



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2016

Formal Total Synthesis of (–)-Jiadifenolide and Synthetic Studies toward seco-Prezizaane-Type Sesquiterpenes

Gomes, José ; Daeppen, Christophe ; Liffert, Raphael ; Roesslein, Joel ; Kaufmann, Elias ; Heikinheimo, Annakaisa ; Neuburger, Markus ; Gademann, Karl

Abstract: Synthetic studies toward highly oxygenated seco-prezizaane sesquiterpenes are reported, which culminated in a formal total synthesis of the neurotrophic agent (–)-jiadifenolide. For the construction of the tricyclic core structure, an unusual intramolecular and diastereoselective Nozaki–Hiyama–Kishi reaction involving a ketone as electrophilic coupling partner was developed. In addition, synthetic approaches toward the related natural product (2R)-hydroxy-norneomajucin, featuring a Mn-mediated radical cyclization for the tricycle assembly and a regioselective OH-directed C–H activation are presented.

DOI: <https://doi.org/10.1021/acs.joc.6b02039>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-134801>

Journal Article

Accepted Version

Originally published at:

Gomes, José; Daeppen, Christophe; Liffert, Raphael; Roesslein, Joel; Kaufmann, Elias; Heikinheimo, Annakaisa; Neuburger, Markus; Gademann, Karl (2016). Formal Total Synthesis of (–)-Jiadifenolide and Synthetic Studies toward seco-Prezizaane-Type Sesquiterpenes. *Journal of Organic Chemistry*, 81(22):11017-11034.

DOI: <https://doi.org/10.1021/acs.joc.6b02039>

**Formal Total Synthesis of (–)-Jiadifenolide and Synthetic Studies Towards
seco-Prezizaane-Type Sesquiterpenes**

José Gomes,[†] Christophe Daeppen,^{†,‡} Raphael Liffert,^{†,‡} Joel Rösslein,^{†,‡} Elias


Kaufmann,^{†,‡} Annakaisa Heikinheimo,[†] Markus Neuburger,[†] and Karl Gademann^{*†,‡}

[†]Department of Chemistry, University of Basel, St. Johannis-Ring 19, CH-4056 Basel,
Switzerland

[‡]Department of Chemistry, University of Zurich, Winterthurerstrasse 190, CH-8057
Zurich, Switzerland

karl.gademann@uzh.ch

TOC Graphic




Abstract

Synthetic studies towards highly oxygenated *seco*-prezizaane sesquiterpenes are reported, which culminated in a formal total synthesis of the neurotrophic agent (–)-jiadifenolide. For the construction of the tricyclic core structure, an unusual intramolecular and diastereoselective Nozaki-Hiyama-Kishi reaction involving a ketone as electrophilic coupling partner was developed. In addition, synthetic approaches towards the related natural product (2*R*)-hydroxynorneomajucin, featuring a Mn-mediated radical cyclization for the tricycle assembly and a regioselective OH-directed C-H activation are presented.

Introduction

Natural products that induce or stimulate neuron differentiation and neurite outgrowth have been identified as promising small molecule alternatives to the nerve growth factor (NGF), a key protein mediating neuritogenesis in humans.¹ Since the early studies of Schreiber, Corey, and Danishefsky, many groups, including ours, have investigated small molecule neurotrophins.² The large variety of chemotypes that are able to stimulate neurite outgrowth suggests chemical interference with different pathways beyond NGF-mediated differentiation and outgrowth. While many pathways remain to be discovered, chemical synthesis might provide a leading role in mapping out these molecular mechanisms of a process central to memory and learning. An interesting class of compounds featuring strong activity as small molecule neurotrophins is represented by the *seco*-prezizaane-sesquiterpenes, in particular those of the majucin-type (Figure 1).³ Several such compounds have been shown to be neurotrophically active,⁴ such as jiadifenin (**1**),⁵ (-)-jiadifenolide (**2**), (2*S*)-hydroxy-3,4-dehydroneomajucin (**3**),⁵ jiadifenoxolane A (**4**)⁶ and the norsesquiterpenoid (2*R*)-hydroxy-norneomajucin (**5**).⁷ Furthermore, also the synthetic carboxylic acid derivative of jiadifenin (**6**) showed potent neurite outgrowth promoting activity in primary cultured rat cortical neurons.^{3b} Biological studies on jiadifenolide (**2**) suggested that the observed neuronal dendritic outgrowth might involve MAP2 AND PSD95 proteins,[Kuzahara, Biochemical and Biophysical Res. Comm 2016, 470, 798] and Shenvi raised the possibility that jiadifenolide might interact with the Cys-loop family of ion channels based on structural similarity to picrotoxinin [Shenvi Nat.Prod.Rep.,2016,33, 535].

Figure 1. Neurotrophic *seco*-prezizaane-sesquiterpenes natural products.




The structurally complex framework of these terpenoids combined with their potent biological activity renders these compounds interesting targets for synthetic studies. A number of successful total syntheses have been reported for both jiadifenin (**1**)⁸ and jiadifenolide (**2**),⁹ culminating in the impressive eight-step, gram-scale synthesis of the latter compound by Shenvi and co-workers.¹⁰ Our research group has been working towards this family of compounds over the past few years,¹¹ and in this publication, we report synthetic studies towards *seco*-prezizaane type sesquiterpenes, and a formal total synthesis of (-)-jiadifenolide (**2**), in full detail.

Results and Discussion

(-)-Jiadifenolide (**2**) is a highly oxygenated sesquiterpene featuring a unique *seco*-prezizaane-type skeleton. Reported strategies for the assembly of the ABC-ring system^{8a-d, 9a-b} involve formation of the rings in the order B→AB→ABC, A→AB→ABC, and C→BC→ABC. So far B-ring cyclization from an AC-ring system is known to proceed *via*


samarium-mediated cross-coupling^{9c-d} or *via* tandem Tsuji-Trost/lactonization reaction.^{8e} We envisioned a novel strategy by building up the B-ring *via* cyclization of the AC-ring system, where the A-ring is originally derived from readily available (*R*)-pulegone *via* methylester **7**¹² and the C-ring from tetronic acid (Scheme 1). For the connection of the A- and C-ring we envisioned a Knoevenagel condensation in order to form the AC-ring fragment.^{2f} The key step, viz. the aforementioned formation of the B-ring, was supposed to be performed by a Nozaki-Hiyama-Kishi (NHK) coupling.¹³ The remaining methyl group at the quaternary stereogenic center C-5 was planned to be installed *via* 1,4-addition to the unsaturated γ -lactone, or a two-step procedure including Corey-Chaykovsky cyclopropanation and subsequent Birch reduction.¹⁴ The bridged D- and E-ring was planned to be constructed by ester reduction and C1-homologation.¹⁵

Scheme 1. Retrosynthetic analysis of (–)-jiadifenolide.



Methylester **7** was synthesized according to a literature procedure¹² in two steps starting from inexpensive (*R*)-pulegone. Subsequent allylation of **7** yielded compound **8** in good yield (92%) and diastereomeric ratio (11:1 in favor of the desired isomer, Scheme 2). Ozonolysis furnished the aldehyde **9**, which was coupled to tetronic acid with Hantzsch ester as reducing agent,¹⁶ yielding the AC-ring fragment **10** in 81% yield. Triflation of the enol **10** led to vinyltriflate **11**, which was treated under NHK reaction conditions.¹³ Pleasingly, diastereoisomer **12** was exclusively formed in 69% yield, and its constitution and configuration was subsequently confirmed by X-ray crystal structure analysis.

Scheme 2. Synthesis of the tetracyclic precursor.




Reagents and conditions: (a) NaH, THF/DMPU (4:1), rt, 1 h, then allyl bromide, 0 °C, 13 h, 92%, dr = 11:1; (b) O₃, CH₂Cl₂/MeOH (5:1), -78 °C; Zn, AcOH, -78 °C to rt, quant.; (c) Tetronic acid, Hantzsch ester, L-proline (10 mol%), MeOH, 24 h, rt, 81%; (d) 2,6-Lutidine, Tf₂O, CH₂Cl₂, -78 °C to rt, 7 h, quant.; (e) CrCl₂, NiCl₂ (10 mol%), 3 Å MS, DMF, 50 °C, 24 h, 69%; (f) BH₃·DMS, THF, 2.5 h, 0 °C to 45 °C, 76%; (g) KCN, THF/H₂O (1:1), rt, 2 d, 65%.

The next challenge was to find an appropriate reducing agent for the transformation of the methylester **12** to the corresponding aldehyde, and to install the D-ring later. The use of LiAlH₄ proved to be too harsh as several products were detected by NMR analysis of the reaction mixture after work-up, whereas reaction with NaBH₄ in ethanol at 60 °C and superhydride (LiEt₃BH) in THF at 0 °C selectively reduced the lactone moiety, while the

methylester remained intact. However, we were very pleased that reduction with borane dimethyl sulfide was found to give aldehyde **13** in 76% yield, and initially expected over reduction to the corresponding diol did not take place. For the introduction of the second γ -lactone, we planned to install the C-11 carbon by cyanide addition followed by hydrolysis and lactonization. Cyanidation with TMS-CN and Et₃N did not lead to the desired product.¹⁷ Potassium cyanide in aqueous THF solution transformed the aldehyde **13** to the tetracyclic dilactone **14** as a single diastereoisomer (65% yield) and the expected cyanohydrin was not observed. The close juxtaposition of the neighboring C-4 hydroxy group to the putative cyanhydrin intermediate could facilitate formation of the corresponding imidate. As the reaction is conducted in aqueous organic solvent, the imine is prone to undergo hydrolysis to form the α -hydroxy lactone **14**. However, the exact mechanism for this observation remains unclear, as multiple other factors might be involved. Regarding the stereoselectivity, diastereoselective nucleophilic attack of the cyanide to the aldehyde would explain the configuration observed. However, also here a mechanism involving both epimeric cyanohydrins can be reasoned, which would be in equilibrium with the aldehyde **13**, and subsequently, only one diastereoisomer would react to the thermodynamic product **14**. Isomerization of the hydroxyl group due to steric effects with the proximate methyl group at C-1 would also be possible, but rather improbable due to the poor acidity at the C-10 position. To the best of our knowledge, this is the first kind of reaction sequence featuring a cyanohydration followed by hydrolysis and γ -lactonization. While this route proved to be effective in installing the ABCD ring system of these natural products, installation of the missing quaternary methyl function on the tetracyclic structure **14** was challenging. 1,4-Addition¹⁸ using various conditions and Corey-Chaykovsky cyclopropanation appeared to be completely unreactive. We assumed that the D-ring lactone in butenolide **14** might be too sterically demanding or influenced the shape of the molecule, rendering a nucleophilic attack at the appropriate position not feasible. In order to test this hypothesis, we applied the conditions evaluated above on the simpler ABC-ring fragment (Scheme 3). Tertiary alcohol **12** was treated with Burgess reagent and the obtained trisubstituted olefin **15** was dihydroxylated to diol **16**. Once again, no reactivity towards cuprate addition¹⁹ as well as Corey-Chaykovsky conditions¹⁴ was observed for both substrates **12** and **16**.

Additionally, Simmons-Smith conditions²⁰ were tested on the diol **16**, but even with the allylic hydroxy group serving as a directing group,²¹ we were not able to cyclopropanate the butenolide moiety.


Scheme 3. Methylation attempts on various substrates.



Reagents and conditions: Burgess reagent, THF, rt, 3.5 h, 93%; (b) OsO₄ (5 mol%), NMO, *t*-BuOH/H₂O (3:1), 0 °C to rt, 3 d, 84%, dr = 10:1.

Sampson and co-workers described a similar scenario and attributed the lack of reactivity to a competitive γ -deprotonation pathway, which affords an aromatic furan oxide rather than the conjugate addition.¹⁴ In order to avoid the reactivity issues for the C-5 methylation we focused on a temporary opening of the A-ring to overcome reactivity issues in this critical methyl group insertion (Scheme 4).

Scheme 4: Synthesis of the aldehyde fragment.




Reagents and conditions: (a) Rh/C/Al₂O₃ (10 mol%), H₂ (60 bar), EtOAc, rt, 3 h, quant.; (b) NaIO₄, THF/H₂O (2:1), rt, 1.5 h, quant.; (c) ethylene glycol, *p*-TSA·H₂O, benzene, 90 °C, 2 h, quant.; (d) formaline, THF, pH 6, 40 °C, 14 h, 94%; (e) PPh₃, imidazole, I₂, benzene, rt, 12 h, 75%; (f) *n*-Bu₃SnH, AIBN, benzene, 85 °C, 3 h, 84%; (g) 1 M HCl, acetone, rt, 14 h, 99%; (h) NaBH(OCH(CF₃)₂)₃, hexafluoro-2-propanol, rt, 16 h, 85%; (i) PPh₃, imidazole, I₂, benzene, rt, 16 h, 95%; (j) SmI₂, HMPA, THF, rt, 1 h, 76%; (k) BH₃:DMS, THF, 2.5 h, 0 °C to 45 °C, 66%.

Olefin hydrogenation on the diol **16** was carried out successfully in quantitative yield in a diastereoselective fashion and the configuration of the hydrogenated product **17** was

confirmed by X-ray crystal structure analysis.²² The A-ring was then cleaved by oxidation to the unstable ketoaldehyde **18** using sodium periodate²³ and protected to obtain the corresponding acetal **19** in quantitative yield. Since methylation of compound **19** did not yield the desired product due to decomposition under basic (MeI with DBU or LDA) and acetal deprotection under acidic conditions, we performed a hydroxymethylation under neutral conditions to install the additional C-5 carbon on substrate **19**. The stereochemical outcome of the reaction was verified by 2D-NMR studies. Hydroxymethylation was expected to be stereospecific due to the formation of the thermodynamically favorable *cis*-fused bicyclic ring system.²⁴ A Garegg-Samuelsson iodination of the OH group in compound **20**, followed by radical dehalogenation of the resulting iodide **21** using tributyltin hydride gave the methylated intermediate **22**. The acetal was hydrolyzed under acidic conditions to form the aldehyde **23**. Reduction of **23** using NaBH₄ was not reproducible, but sodium tris(hexafluoroisopropoxy)borohydride, reported as a very mild reducing reagent,²⁵ was found to be efficient (85%). Primary alcohol **24** was subsequently iodinated applying Garegg-Samuelsson conditions and the structure of the resulting iodide **25** was confirmed by X-ray crystal structure analysis. First attempts to close the A-ring using organolithium reagents resulted in complex mixtures (*n*-BuLi)²⁶ or formation of minor amounts of **26** (*t*-BuLi).^{2c} Finally, SmI₂ mediated ring closing using HMPA as an additive, which was found to be crucial, gave the desired tricycle **26** as a single diastereomer in a remarkable yield of 76%.²⁷ We were pleased that the borane dimethyl sulfide mediated reduction applied earlier on ester **12**, was reproducible for intermediate **26** to yield the desired aldehyde **27** in 66%.

Our initial established strategy for the C1-homologation on the aldehyde **27** was resumed to produce the tetracyclic structure **28** (Scheme 5). Selenoxide promoted elimination²⁸ then formed an α,β -unsaturated intermediate which under the applied reaction conditions cyclized to form the E-ring containing lactone **29**. Oxidation of lactone **29** in the presence of H₂O₂ led directly to pentacycle **33**. Unfortunately, the pentacycle **33** had shown to be completely unreactive towards the final installation of the hydroxyl-group at C-10 using various oxidation protocols²⁹ (Scheme 5). Aware of the difficulties accompanied with the introduction of this critical hydroxyl group, intermediate **29** was treated under different conditions using LiHMDS as base to obtain the α,β -unsaturated lactones **30** or **32**, which should act as stable precursors for the final E-ring formation. However, treating compound **32** under established epoxidation conditions yielded in desilylation and subsequent formation of the aforementioned α -hydroxy lactone **33**.^{4b} Studies using the more stable TES protecting group led to formation of pentacycle **33** as well. It was thus decided to first oxidize the D-ring moiety at C-10 to overcome reactivity issues of intermediate **33**. We oxidized compound **30** with DMP to the corresponding ketone **31**. To our disappointment, the same epoxidation conditions culminated in decomposition of the starting material. Presumably, the electrophilic properties of the ketone dominate over the Michael acceptor capability, which results in decomposition of the starting material.

Scheme 5: Synthetic approaches towards (-)-jiadifenolide




Reagents and conditions: (a) KCN, THF/H₂O, rt, 3 d (63% over two steps); (b) LDA, PhSeBr, H₂O₂, THF, -78 °C to 0 °C, 1.5 h, 48%; (c) LiHMDS, -78 °C to rt, 2 h, n.d.; (d) DMP, CH₂Cl₂, rt, 16 h, n.d.; (e) LiHMDS, TMSCl, THF, -78 °C to 0 °C, 1 h, n.d.; (f) NaOH, H₂O₂, MeOH/THF (3:1), 0 °C to rt, 14 h.

As the introduction of the quaternary methyl group at C-5 proved cumbersome, and was only solved by a lengthy route, we directed our attention to alternative approaches towards the construction for the ABC-ring intermediate along with an earlier installation of this critical methyl group. At first, we aimed to construct the A- and B-ring *via* a


diastereoselective inverse electron-demand Diels-Alder reaction or more likely to use a stepwise approach by the combination of either 1,4- or 1,6-addition and subsequent aldol addition (Scheme 6).

Scheme 6. Retrosynthetic analysis of the AB-ring system.



Different strategies involving aldol cyclisation were evaluated. The 1,4-addition adduct **35** was synthesized by Michael addition of β -ketoester **7** to the literature known³⁰ benzyl protected hydroxy ketone **34** using potassium carbonate in acetone with 70% yield and a diastereoselectivity of 5:1 (Scheme 7). Neither for the benzyl protected alcohol **35** nor the deprotected alcohol **36** an aldol cyclisation took place by using different bases (Cs_2CO_3 , KOH, DBU, NaOMe, *t*-BuOK) or enamine catalysis (pyrrolidine/AcOH, proline). Attempts forming the thermodynamic silyl enol ether suitable for Mukaiyama aldol reaction from protected β -hydroxy ketone **35** (TMSCl, NaI, NEt_3 or TBSOTf, NEt_3) or trapping the formed enolate after reaction of ester **7** with benzyl protected alcohol **34** (LDA, then TBSOTf) were not successful either. The α,β -unsaturated aldehyde **38** was synthesized by 1,6-addition of ester **7** to the known dienal **37**³¹ with K_2CO_3 in acetone at 30°C in 66 % yield.

Scheme 7: Synthesis of the AB-ring system.




Reagents and conditions: a) K_2CO_3 , acetone, 40 °C, o.n., 70%, dr = 5:1; b) 5% Pd/C, EtOH, rt, 2.5 h, quant.; c) K_2CO_3 , acetone, 30 °C, o.n., 66%; d) Me_3Al , LiSPh, THF, 0 °C to 40 °C, 37%, dr = 9:5.

Cyclization of **38** via the enamine did not take place, most likely due to the challenges in the isomerization of the conjugated double bond. Nevertheless, the enamine of **38** (piperidine, AcOH, 105 °C) could be detected via ^1H -NMR spectroscopy. In order to circumvent the problems accompanied by this double bond we considered using a temporary protecting group for this critical functional group. Aldol reaction of aluminium enolates generated by 1,4-addition of organoaluminium or organoselenium reagents to α,β -unsaturated carbonyl compounds was described by Nozaki.³² Following the

cyclization protocol applied for the oxahydrindene fragment from Danishefsky's synthesis of Avermectin A_{1a},³³ we treated aldehyde **38** with the species formed from the reaction of trimethylaluminium and lithium thiophenolate. Instead of the desired AB-ring product, Michael addition to form a diastereomeric mixture (dr = 9:5) of the bicyclo[3.2.1]octanone **39** in 37 % yield was observed.

Scheme 8: Hosomi-Sakurai approach.




Reagents and conditions: a) acrolein, NEt₃, CH₂Cl₂, rt, o.n., 71%; b) isopropenylmagnesium bromide, THF, -78 °C, 3 h, 33%; c) vinylmagnesium bromide, THF, -78 °C, 4 h, 42%; d) DMP, CH₂Cl₂, rt, 1 h, quant.; e) DMP, CH₂Cl₂, rt, 2 h, quant.; f) allyltrimethylsilane, Grubb's II (5 mol%), CH₂Cl₂, 40 °C, 90 min, 71%; g) allyltrimethylsilane, Grubb's II (5 mol%), CH₂Cl₂, 40 °C, 90 min, 51%; h) PivCl, pyridine, DMAP, CH₂Cl₂, rt to 40 °C, o.n. crude; i) BnO(NH)CCl₃, Sc(OTf)₃ (5 mol%), CH₂Cl₂, 0 °C, 90 min, 88%.

Due to the failure of the cyclization of the B-ring using aldol addition protocols, we planned to use a diastereoselective intramolecular Hosomi-Sakurai reaction³⁴ instead (Scheme 8). Different strategies to efficiently prepare the allylsilane precursor were

evaluated. Attempts to install the TMS group at C-7 using **7** and 2-(trimethylsilyl)acrylaldehyde or *via* introduction of the allylsilane part by Grubbs metathesis at the C-5 position proved to be unsuccessful. Finally, we were pleased to find a route to synthesize test substrates suitable for the Hosomi-Sakurai reaction but with a missing methyl group. The synthesis of these substrates started by Michael-addition of **7** to acrolein, followed by Grignard reaction of isopropenyl- or vinylmagnesium bromide to the formed aldehyde **40** to yield **41** and **42**, respectively. Cross metathesis using Grubbs 2nd generation catalyst gave a moderate yield (51%) for the vinyl substrate **42** bearing the free alcohol. Higher yields (71%) were observed for the Dess-Martin periodinane oxidized substrate **45**, however for PMB protected substrate **43** no metathesis product was isolated. Neither the free alcohol **41**, nor the oxidized substrate **44**, derived from the reaction of **40** with isopropenyl Grignard reagent, were reactive in the metathesis reaction and only homocoupling of allyltrimethylsilane was observed using different reaction conditions. The Hosomi-Sakurai reaction was probed on the test substrates **46**, **47** and **48**, but even without bearing the sterically demanding methyl group at C-5 no reaction took place.

Scheme 9. Formal synthesis of (–)-jiadifenolide



Reagents and conditions: (a) MeMgBr, Fe(acac)₃, NMP, THF, -30 °C, 1 h, 96%; (b) SmI₂, THF/H₂O, rt, 13 h, 50%, dr = 5:2; (c) BH₃·DMS, THF, 6 h, 0 °C to 45 °C, 51%, dr = 5:1.


Zhang and co-workers recently published the protecting group free total synthesis of (-)-jiadifenolide (**2**).^{9d} Their synthetic route lead through the same aldehyde intermediate **27**, which was constructed *via* a SmI₂ mediated closure of the B-ring and formed directly the critical methyl function bearing quaternary C-5 stereocenter. Radical cyclization of the B-ring of substrate **49** was investigated in our group as well, as the approaches to install the methyl group on the C ring proved cumbersome. β-Methylated lactone **49** was synthesized by applying Fürstner's conditions to vinyltriflate **11** (Scheme 9). To our disappointment, initial evaluation of standard conditions did not facilitate the desired cyclization of **49**. Reaction with SmI₂ in THF without any further additives led mostly to the formation of an unidentified side-product at 0 °C as well as at reflux. Additives such as HMPA and NiI₂ or the addition of protic additives such as MeOH and *t*-BuOH did not lead to the formation of the desired product **26**. Given the encouraging results reported by Zhang and coworkers on the addition of excess water being crucial for the SmI₂-mediated radical cyclization, we tested these conditions on the substrate **49**. After small adaptations, we were able to isolate the desired cyclized product **26** in 50% yield and a diastereomeric

ratio of 5:2 (not separable) for C-5. Additionally, a side-product where the keto group was reduced to the OH function³⁵ could be isolated in 39% yield (dr 13:1). Treating the diastereomeric mixture of **26** with BH₃DMS furnished intermediate **27** in a diastereomeric ratio of 5:1 (51% isolated yield) along with recovered starting material (15%) in a diastereomeric ratio of 4:3. Concerning the increase of the diastereomeric ratio in the transformation from **26** to **27**, we suggest that in the minor diastereomer of **26** the methyl group at C-5 *syn* to the ester residue has a steric influence and slows down the reduction to the aldehyde. In contrast, a coordination of the butenolide moiety in the major diastereomer of **26** to the BH₃ might support the reduction to aldehyde **27**. In summary, aldehyde **27** was obtained from (*R*)-pulegone in a nine step procedure and an overall yield of 10%.

To gain access to a variety of majucin-type sesquiterpenes *via* derivatization, we targeted on a different synthetic approach with involving a similar precursor also represented by a common ABC-ring fragment. Manganese induced radical cyclization, already applied in our group and successfully utilized in the synthesis of endoperoxides,³⁶ might constitute a feasible approach for the construction of the B-ring. Functionalization of the geminal methyl functions is present in broad range of *seco*-prezizaane-type sesquiterpenes. The known tetrasubstituted olefin **50** would be an ideal substrate for a regiospecific sp³ C-H oxidation of the geminal methyl group, which will be incorporated in the butenolide moiety.

Starting from enantiopure (*R*)-pulegone, the known intermediate **50** was readily accessible (Scheme 11).¹² α -Alkylation attempts using LiHMDS or LDA without additives were found to only deprotonate one of the diastereomers, but addition of DMPU triggered the reaction to full conversion and the desired alkylated ester **51** was obtained as a single diastereomer. The TBS-protecting group was cleaved with TBAF/AcOH and the formed alcohol **52** was further oxidized with DMP to the desired aldehyde **53** in 89%. A Roskamp reaction³⁷ furnished the desired β -ketoester **54** in an excellent yield of 93%. Initial screening for the manganese induced radical cyclization showed formation of the desired B-ring. However, the cyclized product **55** was isolated in low yield and along with side-products. Raising the temperature to 50 °C could reduce the reaction time but byproduct formation was not eliminated. Slow addition of the Mn(OAc)₃ or of the substrate had no influence and an average yield of 25-30%, with 39% as the best result was obtained. The byproducts were only isolated as complex mixtures, suggesting that several intermolecular radical recombination reactions occurred. Instead of the expected disubstituted compound, the trisubstituted olefin **55** was directly isolated, which might result from chloride elimination. Ketone **55** was reduced in the presence of NaBH₄ in a complete diastereoselective manner and alcohol **56** could be secured by X-ray crystallography analysis. With the desired hydroxy function in hand, a regioselective C-H functionalization to form the ether **57** was achieved in the presence of Pb(OAc)₄ in 81% yield.³⁸ The byproduct **58**, bearing an additional acetyl group at the carbon next to the ether,³⁹ was further isolated and successfully converted to the desired product **57** using BF₃·OEt₂ and triethylsilane.⁴⁰ Both structures were confirmed by X-ray crystallographic analysis (Scheme 11).

Scheme 11. Manganese-mediated cross-reaction to establish the AB-core




Reagents and conditions: (a) LDA, DMPU, THF, *tert*-butyl(2-iodoethoxy)dimethylsilane,⁴¹ -78°C to rt, 77%; (b) TBAF, AcOH, CH_2Cl_2 , rt, 16 h, 79%; (c) DMP, CH_2Cl_2 , rt 4 h, 89%; (d) ethyl 2-diazoacetate, SnCl_2 (cat.), CH_2Cl_2 , 1 h, 93%; (e) $\text{Mn}(\text{OAc})_3$, LiCl, AcOH, 50°C , 5 h, 39%; (f) NaBH_4 , MeOH, 0°C to rt, 4 h, 86%; (g) $\text{Pb}(\text{OAc})_4$, cyclohexane, I_2 , CaCO_3 , hv, 3 d, 81%; (h) $\text{BF}_3\cdot\text{OEt}_2$, Et_3SiH , CH_2Cl_2 , -78°C to rt, 15 h, 88%.

To our surprise, the ether bridge was found to be stable under several conditions.⁴²

Therefore, the chloride **57** was reduced by tributyltin hydride to form the ether **59** in 95% yield (Scheme 12). The bridged ether was then opened in the presence of Ac_2O and BF_3 -etherate⁴³ to give the diacetate **60** along with the elimination byproduct **61**, and the acetyl groups of **60** were cleaved with K_2CO_3 in MeOH to form the corresponding diol **62**. The configuration of the OH group could not be assigned in compound **62**, and subsequent TPAP-mediated oxidation to the enolaldehyde **63** was carried out. Other oxidation conditions furnished enolaldehyde **63** as well, but were found to be not reproducible. Conditions using IBX showed traces of the overoxidized α -hydroxylated product **64**, which indicated a general accessibility of the missing OH function insertion at this stage. Nevertheless, the method of choice for the preparation of **64** turned out to be exposure of

63 to CeCl_3 and molecular oxygen to yield the α -hydroxylated product **64** in 43%.⁴⁴ A reductive cyclization of **64** formed the butenolide and also caused partial closure of the D-ring. This mixture was subjected to acid mediated lactonization involving *p*-TsOH to fully promote D-ring formation in compound **65**, however in a very low yield of 27% over the two steps. Final modifications of the A-ring are proposed in order to give access to the natural product (2*R*)-hydroxynorneomajucin (**5**) from tetracycle **65**. This intermediate **65** might also serve as a common precursor to other *nor*-type *seco*-prezizaane sesquiterpenes.

Scheme 12. Synthesis of the (2*R*)-hydroxy-norneomajucin precursor.



Reagents and conditions: (a) Bu_3SnH , AIBN, benzene, 100 °C, 2 h, 95%; (b) $\text{BF}_3 \cdot \text{OEt}_2$, AcOH, -20 °C to rt, 12 h, 60%; (c) K_2CO_3 , MeOH, rt, 16 h, 98%; (d) TPAP, NMO, 4 Å MS, CH_2Cl_2 , RT, 16 h, 68%; (e) $\text{CeCl}_3 \times 7 \text{ H}_2\text{O}$, O_2 , *i*-PrOH, rt, 24 h, 43%; (f) NaBH_4 , MeOH, 0 °C to rt, 12 h, then *p*-TsOH, PhMe, 70 °C, μw , 2 h, 27% (2 steps).

Conclusion

In conclusion, we present a new strategy to access the tricyclic core structure of (–)-jiadifenolide *via* a Nozaki-Hiyama-Kishi coupling, which set the stage for an diastereoselective, formal total synthesis of the natural product. Due to the challenging introduction of the methyl group in order to construct this quaternary stereogenic center after the B-ring closure, we elaborated a more efficient route to the tricycle by using a SmI₂-mediated radical cyclization. Unfortunately, alternative strategies to close the B-ring using aldol-cyclization protocols or a Hosomi-Sakurai reaction were unsuccessful. Furthermore, a C1-homologation promoted by potassium cyanide successfully provided the D-ring formation in a remarkable one-step procedure to allow for the completion of the synthesis *via* successive oxidation protocols. Synthetic studies towards (2*R*)-hydroxynorneomajucin were carried out, which involved a manganese mediated radical cyclization and a regiospecific C-H activation. Moreover, these studies potentially allow to access a variety of *nor*-derivative sesquiterpene natural products as potential neurotrophic modulators.

Experimental Section

General, Materials and Equipment

All chemicals have been purchased from Acros, Fluka or Sigma-Aldrich and were used without further purification (except for Et₃N which was freshly distilled before use). All reactions have been carried out in heat gun-dried glassware (unless aqueous reagents

were used) and reactions involving air sensitive compounds have been performed under an argon or nitrogen atmosphere. Solvents applied for chemical transformations were either puriss quality or HPLC grade solvents, which have been dried by filtration through activated aluminium oxide under nitrogen (H_2O content <10 ppm, *Karl-Fischer* titration). For work-up and purification solvents have been distilled from technical grade. All synthetic transformations have been monitored by either thin layer chromatography (TLC) or ^1H NMR spectroscopy. Yields refer to purified, dried and spectroscopically pure compounds. TLC was performed on Merck silica gel 60 F254 plates (0.25 mm thickness) precoated with fluorescent indicator. Concentration under reduced pressure was performed by rotary evaporation at 40°C . Flash chromatography was performed using silica gel 60 (230-400 mesh) from SiliCycle or Sigma-Aldrich with a forced flow eluent at 0.2-0.4 bar pressure. All ^1H and ^{13}C NMR spectra were recorded using a Bruker Avance 250 MHz, 400 MHz or 500 MHz (^1H) & 63 MHz, 101 MHz or 126 MHz (^{13}C) & 376 MHz (^{19}F) spectrometer at room temperature. Chemical shifts (δ -values) are reported in ppm, spectra were calibrated related to solvents' residual proton chemical shifts (CHCl_3 , $\delta = 7.26$; $\text{d}_4\text{-MeOD}$, $\delta = 3.31$) and solvents' residual carbon chemical shifts (CDCl_3 , $\delta = 77.16$; $\text{d}_4\text{-MeOD}$, $\delta = 49.00$), multiplicity is reported as follows: s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet or unresolved and coupling constant J in Hz. IR spectra were recorded on a Varian 800 FT-IR ATR Spectrophotometer or a Perkin Elmer SpectrumTwo ATR-FTIR. The absorptions are reported in cm^{-1} . All masses spectra (HRMS-ESI) were recorded by the Mass spectrometric Service of the University of Bern on Sciex QSTAR Pulsar mass spectrometer using electrospray ionization, University of Basel on a QTOF-ESI Spectrometer (bruker maXis 4G) or University of Zürich using a high-resolution mass spectra acquired on a QExactive instrument (Thermo Fisher Scientific, Bremen, Germany) equipped with a heated electrospray (ESI) ionization source and connected to a Dionex Ultimate 3000 UHPLC system. Melting points (M.p.) were determined using a Büchi B-545 apparatus in open capillaries and are uncorrected. Optical rotations $[\alpha]_D^T$ were measured at the sodium D line using a 1 mL cell with a 1 dm path length on a Jasco P-2000 digital polarimeter and the concentration c is given in g/100 mL and the used solvent is CHCl_3 . Elementary analysis were performed at the University of Basel. X-ray analysis: Data collection for the crystal structure was

performed at low temperature using Mo K_{α} radiation on a Bruker KappaAPEX diffractometer. Integration of the frames and data reduction was carried out using APEX2.⁴⁵ The structures were solved by direct methods using SIR92.⁴⁶ All non-hydrogen atoms were refined by full-matrix least-squares on F using CRYSTALS.⁴⁷ Hydrogen atoms were placed in calculated positions by means of the “riding” model.

Methyl (2R)-2-methyl-5-oxocyclopentane-1-carboxylate (7): Methyl (2R)-2-methyl-5-(propan-2-ylidene)cyclopentane-1-carboxylate²¹ (360 mg, 1.98 mmol, 1 eq.) was dissolved in MeOH (20 mL) and CH_2Cl_2 (100 mL) at $-78\text{ }^{\circ}\text{C}$. Ozone was bubbled in the solution until the solution became light blue. Ozone addition was stopped and the solution was stirred under argon at $-78\text{ }^{\circ}\text{C}$ for 15 min and at rt for 10 min. Zinc powder (526 mg, 7.96 mmol, 4.0 eq.) and acetic acid (0.46 mL, 7.96 mmol, 4.0 eq.) were added to the solution. The resulting mixture was stirred at rt for 30 min and filtered over celite. The filtrate was neutralized with saturated NaHCO_3 solution and the aqueous layer was extracted two times with CH_2Cl_2 . The combined organic layers were washed with water and brine, dried over Na_2SO_4 , filtered and evaporated under reduced pressure to afford ketone **7** (300.6 mg, 1.92 mmol, 97%) as colorless oil: $R_f = 0.13$ (pentane/ Et_2O 9:1); $[\alpha]_D^{26} = +93.5$ (c 0.46, CHCl_3). FTIR (neat); $\tilde{\nu} = 2959, 2875, 1754, 1724, 1436, 1335, 1289, 1260, 1202, 1128, 1006\text{ cm}^{-1}$; ^1H NMR (400 MHz, CDCl_3) $\delta = 3.76$ (s, 3H), 2.77 (d, $J = 11.4\text{ Hz}$, 1H), 2.67–2.53 (m, 1H), 2.48–2.39 (m, 1H), 2.38–2.26 (m, 1H), 2.20 (dddd, $J = 12.7, 8.3, 6.2, 2.0\text{ Hz}$, 1H), 1.48 (dtd, $J = 12.6, 11.2, 8.5\text{ Hz}$, 1H), 1.18 (d, $J = 6.5\text{ Hz}$, 3H); ^{13}C NMR (101 MHz, CDCl_3) $\delta = 212.0, 169.7, 63.1, 52.5, 38.9, 36.5, 29.5, 19.4$; elemental analysis calcd for $\text{C}_8\text{H}_{12}\text{O}_3$: C = 61.52; H = 7.74; N = 0.00; found: C = 61.49; H = 7.60; N = 0.00; HRMS (ESI) m/z calcd for $\text{C}_8\text{H}_{13}\text{O}_3$ $[\text{M} + \text{H}]^+$: 157.0859, found: 157.0861.

Methyl (1R,2R)-1-allyl-2-methyl-5-oxocyclopentane-1-carboxylate (8): Ketone **7** (1.00 g, 6.4 mmol, 1.0 eq.) was dissolved in dry THF (14.0 mL) and freshly distilled DMPU (3.6 mL). Sodium hydride (60% dispersion in mineral oil, 307 mg, 7.7 mmol, 1.2 eq.) was added at rt and the grey suspension was stirred for 1 h after which it turned to a yellow

solution. Allyl bromide (0.67 mL, 7.7 mmol, 1.2 eq.) was added at 0 °C and the milky mixture was stirred for 1 h at 0 °C before it was allowed to warm up to rt. After 12 h the mixture was diluted with Et₂O (20 mL) and washed three times with water. The aqueous layers were re-extracted twice with Et₂O. The combined organic layers were dried over Na₂SO₄, filtered and evaporated under reduced pressure. Both diastereoisomers were separated by chromatography (pentane/Et₂O 9:1) giving the desired allylated compound **8** (1.06 g, 5.4 mmol, 84%) as colorless oil; *R_f* = 0.48 (pentane/Et₂O 4:1); [α]_D²⁵ = +38.4 (*c* 0.25, MeOH); FTIR (neat): $\tilde{\nu}$ = 3079, 2958, 2882, 1749, 1731, 1434, 1227, 1187, 1165, 998, 919 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ = 5.56–5.67 (m, 1H), 5.09–5.13 (m, 1H), 5.08 (s, 1H), 3.69 (s, 3H), 2.66 (dd, *J* = 14.2, 6.2 Hz, 1H), 2.47–2.58 (m, 2H), 2.27–2.38 (m, 1H), 2.00–2.18 (m, 2H), 1.72–1.84 (m, 1H), 1.02 (d, *J* = 6.9 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ = 216.2, 171.1, 133.2, 119.6, 63.1, 52.0, 39.1, 38.7, 35.8, 28.3, 15.6; elemental analysis calcd for C₁₁H₁₆O₃: C 67.32; H = 8.22; found: C 67.29; H = 8.20; EI-MS 70 eV, *m/z* (%) = 196.1 (22, M⁺), 168.1 (100), 136.1 (63), 109.1 (90), 81.1 (68).

Methyl (1R,2R)-2-methyl-5-oxo-1-(2-oxoethyl)cyclopentane-1-carboxylate (9): Alkene **8** (2.50 g, 12.8 mmol, 1.0 eq.) was dissolved in a CH₂Cl₂/MeOH mixture (30 mL, 5:1) and cooled down to –78 °C. Ozone was bubbled through the stirring mixture until it turned slightly blue. The addition of ozone was stopped and the solution was stirred for 15 minutes at –78 °C during which time the mixture turned colorless again. Zinc powder (3.36 g, 51.0 mmol, 4.0 eq.) and acetic acid (2.6 mL, 51.0 mmol, 4.0 eq.) were added. The resulting mixture was allowed to warm up to rt overnight. The mixture was filtered over celite and washed with CH₂Cl₂. After evaporation of the solvent, the residue was suspended with CH₂Cl₂, filtered, and evaporated again to give aldehyde **9** (2.54 g, 12.8 mmol, quant.) as colorless oil; *R_f* = 0.68 (pentane/Et₂O 9:1); [α]_D²⁵ = +16.4 (*c* 0.51, MeOH); FTIR (neat): $\tilde{\nu}$ = 2959, 2881, 2857, 2746, 1749, 1717, 1629, 1599, 1456, 1435, 1401, 1397, 1350, 1326, 1233, 1217, 1197, 1167, 1126, 1059, 1038, 1006, 904, 759 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ = 9.66 (d, *J* = 0.5 Hz, 1H), 3.71 (s, 3H), 3.13 (d, *J* = 18.8 Hz, 1H), 2.91 (dd, *J* = 18.8, 1.2 Hz, 1H), 2.54–2.60 (m, 2H), 2.36–2.47 (m, 1H), 2.08–2.18 (m, 1H), 1.79–1.92 (m, 1H), 1.01 (d, *J* = 6.9 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ = 215.3, 199.3, 170.3, 60.5, 52.4, 46.1, 40.1, 38.0, 28.6, 15.7; HRMS (ESI) *m/z*

calcd for C₁₀H₁₅O₄ [M + H]⁺: 199.0965, found: 199.0963.

Methyl (1R,2R)-1-(2-(4-hydroxy-2-oxo-2,5-dihydrofuran-3-yl)ethyl)-2-methyl-5-oxocyclopentane-1-carboxylate (10): A solution containing the aldehyde **9** (0.49 g, 2.47 mmol, 1.0 eq.), Hantzsch ester (1.38 g, 5.44 mmol, 2.2 eq.), tetronic acid (0.26 g, 2.47 mmol, 1.0 eq.) and L-proline (28.5 mg, 0.25 mmol, 0.1 eq.) in methanol (30 mL) was stirred for 24 h at rt. After evaporation under reduced pressure, the residue was purified by flash chromatography (CH₂Cl₂/MeOH 20:1) to give ketone **10** (0.56 g, 1.99 mmol, 81%) as a beige solid; Mp: 125.5–125.9 °C; R_f = 0.23 (CH₂Cl₂/MeOH 10:1); [α]_D²⁵ = +12.8 (c 0.27, MeOH); FTIR (neat): $\tilde{\nu}$ = 2961, 2942, 2882, 2699, 1746, 1729, 1714, 1648, 1406, 1231, 1195, 1104, 1041, 988, 767 cm⁻¹; ¹H NMR (400 MHz, d₄-MeOD) δ = 4.57 (s, 2H), 3.67 (s, 3H), 2.41–2.52 (m, 2H), 2.23–2.34 (m, 2H), 2.05–2.16 (m, 2H), 1.85–2.17 (m, 2H), 1.77 (ddd, *J* = 23.8, 11.5, 8.5 Hz, 1H), 1.05 (d, *J* = 6.9 Hz, 3H); ¹³C NMR (126 MHz, d₄-MeOD) δ = 218.3, 178.4, 175.6, 172.6, 100.8, 68.3, 64.1, 52.2, 40.7, 39.7, 30.6, 29.3, 17.0, 16.1; elemental analysis calcd for C₁₄H₁₈O₆: C = 59.57; H = 6.43; found: C = 59.27; H = 6.20; EI-MS 70 eV, *m/z* (%) = 282.1 (1, M⁺), 251.1 (9), 157.2 (77), 141.1 (100), 109.1 (40).

Methyl (1R,2R)-2-methyl-5-oxo-1-(2-(2-oxo-4-(((trifluoromethyl)sulfonyl)oxy)-2,5-dihydrofuran-3-yl)ethyl)cyclopentane-1-carboxylate (11): Enol **10** (228 mg, 0.81 mmol, 1.0 eq.) and 2,6-lutidine (95 μl, 0.81 mmol, 1.0 eq.) were dissolved in dry CH₂Cl₂ (10 mL) and cooled down to –78 °C. Trifluoromethanesulfonic anhydride (0.14 mL, 0.81 mmol, 1.0 eq.) was added dropwise in 20 minutes. The reaction mixture was allowed to warm up to rt over 7 h. The yellow solution was treated with pentane (10 mL), filtered and the filtrate was extracted with EtOAc, washed with saturated NaHCO₃ solution and brine. The organic layers were dried over Na₂SO₄, filtered and evaporated. Purification by flash chromatography (EtOAc/pentane 1:4) gave vinyltriflate **11** (334 mg, 0.81 mmol, quant.) as a slightly yellow solid; Mp: 43.6–44.4 °C; R_f = 0.33 (EtOAc/pentane 1:4); [α]_D²⁶ = +7.2 (c 0.28, MeOH); FTIR (neat): $\tilde{\nu}$ = 2962, 2885, 2361, 2340, 1778, 1750, 1731, 1703, 1434, 1221, 1136, 1087, 1046, 816 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ = 4.90–4.92 (m, 2H), 3.71 (s, 3H), 2.72–2.52 (m, 2H), 2.41–2.20 (m, 3H), 2.01–2.14 (m,

2H), 1.73–1.95 (m, 2H), 1.07 (d, $J = 6.8$ Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) $\delta = 215.7$, 170.5, 169.8, 160.5, 120.3, 66.6, 62.2, 52.1, 41.2, 38.6, 29.4, 28.3, 17.8, 16.0; ^{19}F NMR (376 MHz, CDCl_3) $\delta = -72.81$; elemental analysis calcd for $\text{C}_{15}\text{H}_{17}\text{F}_3\text{O}_8\text{S}$: C = 43.48; H = 4.14; found: C = 43.60; H = 3.95; EI-MS 70 eV, m/z (%) = 281.1 (16), 249.1 (52), 221.1 (57), 166.2 (62), 141.1 (100), 109.1 (68).

Methyl (5aR,6R,8aR)-8a-hydroxy-6-methyl-3-oxo-1,3,4,5,6,7,8,8a-octahydro-5aH-indeno[4,5-c]furan-5a-carboxylate (12): Nickel(II) chloride (106 mg, 0.82 mmol, 0.1 eq.), chromium(II) chloride (3.12 g, 7.72 mmol, 6.0 eq.) and some 3 Å molecular sieves were dried in a round bottom flask under reduced pressure with a heat gun. Triflate **11** (1.70 g, 4.11 mmol, 1.0 eq.) dissolved in dry DMF (40 mL) was degassed and subsequently added to the dried chromium- and nickel-mixture. The green mixture was stirred at 50 °C for 24 h, then quenched by water (100 mL) and extracted with Et_2O until no more product could be detected in the aqueous layer by TLC. The combined organic layers were dried over Na_2SO_4 , filtered and evaporated. The residue was purified by flash chromatography (EtOAc /pentane 1:4) to afford the cyclized product **12** (749.0 mg, 2.81 mmol, 69%) as a white solid; Mp: 129.5–129.7 °C; $R_f = 0.30$ (EtOAc /pentane 1:3); $[\alpha]_D^{25} = -37.0$ (c 0.26, MeOH); FTIR (neat): $\tilde{\nu} = 3346$, 2956, 2899, 2360, 1728, 1675, 1443, 1245, 1194, 1153, 1100, 1058, 1015, 653 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) $\delta = 5.06$ (d, $J = 1.9$ Hz, 1H), 4.96–5.03 (m, 1H), 4.78 (dt, $J = 17.5$, 2.8 Hz, 1H), 3.77 (s, 3H), 2.25–2.36 (m, 2H), 2.01–2.24 (m, 4H), 1.85–1.95 (m, 2H), 1.73–1.83 (m, 1H), 1.11 (d, $J = 6.8$ Hz, 3H); ^{13}C NMR (126 MHz, CDCl_3) $\delta = 176.5$, 173.4, 164.7, 124.8, 80.9, 64.9, 60.2, 52.4, 41.7, 38.5, 32.9, 31.2, 20.4, 18.7; elemental analysis calcd for $\text{C}_{14}\text{H}_{18}\text{O}_5$: C = 63.15; H = 6.81; found: C = 63.02; H = 6.80; EI-MS 70 eV, m/z (%) = 266.1 (45, M^+), 238.1 (31), 206.1 (82), 178.1 (100), 133.1 (39), 91.1 (26); X-ray crystal structure is given in the supporting information.

(5aR,6R,8aR)-8a-Hydroxy-6-methyl-3-oxo-1,3,4,5,6,7,8,8a-octahydro-5aH-indeno[4,5-c]furan-5a-carbaldehyde (13): Borane dimethyl sulfide complex (2 M in THF, 0.83 mL, 1.65 mmol, 2.0 eq.) was added dropwise to a solution of the methylester **12** (220 mg, 0.83 mmol, 1.0 eq.) in dry THF (4 mL) at 0 °C. The colorless solution was stirred

subsequently for 0.5 h and then warmed up to 45 °C for 2 h. After cooling down to rt, ethanol was added until the exothermic reaction and gas evolution ceased. After evaporation, the residue was purified by flash chromatography (Et₂O/pentane 3:1) giving aldehyde **13** (149 mg, 0.63 mmol, 76%) as a white solid; Mp: 158.1–158.5 °C; *R_f* = 0.21 (Et₂O/pentane 3:1); $[\alpha]_D^{26} = -49.3$ (*c* 0.23, MeOH); FTIR (neat): $\tilde{\nu} = 3346, 2942, 2874, 2758, 1744, 1716, 1669, 1435, 1353, 1224, 1107, 1061, 1016, 929, 753\text{ cm}^{-1}$; ¹H NMR (400 MHz, CDCl₃) $\delta = 9.70$ (s, 1H), 5.01 (ddd, *J* = 17.4, 3.4, 1.7 Hz, 1H), 4.80 (dt, *J* = 17.4, 2.7 Hz, 1H), 4.07 (d, *J* = 1.7 Hz, 1H), 2.04–2.36 (m, 6H), 1.72–1.99 (m, 3H), 1.17 (d, *J* = 7.1 Hz, 3H); ¹³C NMR (126 MHz, CDCl₃) $\delta = 206.8, 173.1, 163.5, 125.2, 80.1, 69.5, 61.9, 40.3, 37.7, 31.6, 28.6, 18.8, 17.7$; elemental analysis calcd for C₁₃H₁₆O₄: C = 66.09; H = 6.83; found: C = 65.90; H = 6.75; EI-MS 70 eV, *m/z* (%) = 236.1 (13, M⁺), 190.1 (100), 175.1 (66), 145.1 (42), 91.1 (27).

(5*aS*,6*R*,8*aR*,11*S*)-11-Hydroxy-6-methyl-4,5,7,8-tetrahydro-6*H*-8*a*,5*a*-(epoxyethano)indeno[4,5-*c*]furan-3,10(1*H*)-dione (**14**): To the aldehyde **13** (43.0 mg, 0.18 mmol, 1.0 eq.), dissolved in THF/water (1:1, 5 mL), was added potassium cyanide (42.0 mg, 0.64 mmol, 3.5 eq.). The resulting mixture was stirred for 2 d at rt before the aqueous reaction mixture was extracted several times with EtOAc until no more product could be detected in the organic layer by TLC. The combined organic layers were dried over Na₂SO₄, filtered and evaporated. The crude product was purified by flash chromatography (EtOAc/cyclohexane 1:1) giving tetracyclic compound **14** (31.0 mg, 0.12 mmol, 65%) as a white solid; Mp: 194.0–194.3 °C; *R_f* = 0.28 (EtOAc/cyclohexane 1:1); $[\alpha]_D^{25} = -32.6$ (*c* 0.19, MeOH); FTIR (neat): $\tilde{\nu} = 3479, 2953, 2929, 2874, 2361, 1785, 1725, 1682, 1442, 1359, 1217, 1191, 1133, 1086, 1046, 1022, 967\text{ cm}^{-1}$; ¹H NMR (400 MHz, CDCl₃) $\delta = 4.82\text{--}4.99$ (m, 2H), 4.57 (d, *J* = 2.8 Hz, 1H), 2.91 (d, *J* = 3.2 Hz, 1H), 2.45–2.54 (m, 2H), 2.21–2.30 (m, 1H), 2.00–2.10 (m, 3H), 1.83–1.98 (m, 2H), 1.62–1.70 (m, 1H), 1.42–1.53 (m, 1H), 1.16 (d, *J* = 6.7 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) $\delta = 176.3, 172.5, 157.5, 127.8, 87.4, 69.6, 68.9, 56.2, 36.2, 35.9, 30.7, 21.3, 15.8, 14.0$; HRMS (ESI) *m/z* calcd for C₁₄H₁₇O₅ [M + H]⁺: 265.1071, found: 265.1065; X-ray crystal structure is given in the supporting information.

Methyl (5aR,6R)-6-methyl-3-oxo-1,3,4,5,6,7-hexahydro-5aH-indeno[4,5-c]furan-5a-carboxylate (15): Ester **12** (301 mg, 1.13 mmol, 1.0 eq.) was dissolved in dry THF (16.0 mL) and Burgess reagent (417 mg, 1.70 mmol, 1.5 eq.) was added. The clear solution was stirred for 3.5 h at rt. Water (8.0 mL) was added and the aqueous layer extracted with Et₂O (3 x 25.0 mL). The combined organic layers were washed with brine (13.0 mL) and then dried over Na₂SO₄, filtered and evaporated to obtain a white solid. The residue was purified by flash chromatography (pentane/EtOAc 3:1) to obtain the desired olefin **15** (262 mg, 1.06 mmol, 93%) as a white solid; Mp: 124.9–125.6 °C; *R_f* = 0.24 (pentane/EtOAc 4:1); $[\alpha]_D^{25} = +193.5$ (*c* 0.16, CHCl₃); FTIR (neat) $\tilde{\nu} = 2952, 2936, 2846, 1740, 1715, 1644, 1439, 1346, 1239, 1172, 1042, 1014, 980, 761, 744 \text{ cm}^{-1}$; ¹H NMR (400 MHz, CDCl₃) $\delta = 6.18$ (s, 1H), 4.98–4.81 (m, 2H), 3.67 (s, 3H), 2.74–2.56 (m, 1H), 2.52–2.15 (m, 2H), 1.35 (ddd, *J* = 12.8, 11.7, 5.8 Hz, 1H), 1.07 (d, *J* = 7.0 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) $\delta = 173.8, 173.1, 151.8, 136.2, 133.2, 126.5, 69.1, 59.9, 52.0, 45.8, 40.6, 30.8, 19.7, 14.9$; HRMS (ESI) *m/z* calcd for C₁₄H₁₇O₄ [*M* + *H*]⁺: 249.1121, found: 249.1119; X-ray crystal structure is given in the supporting information.

Methyl (5aR,6R,8R,8aS)-8,8a-dihydroxy-6-methyl-3-oxo-1,3,4,5,6,7,8,8a-octahydro-5aH-indeno[4,5-c]furan-5a-carboxylate (16): Olefin **15** (80.0 mg, 0.32 mmol, 1.0 eq.) and NMO (49.7 mg, 0.48 mmol, 1.5 eq.) were dissolved in *tert*-butanol (3 mL) and water (1 mL). The clear solution was cooled to 0° C and OsO₄ (4% solution in water, 0.13 mL, 16.1 μmol, 5 mol%) was added dropwise. The clear solution was then allowed to warm up to rt and then stirred for 3 days. To the yellow solution was added saturated Na₂SO₃ solution (15 mL) and the orange to yellow emulsion was stirred for 30 minutes at rt. The mixture was extracted with EtOAc (3 x 10 mL) and the combined organic layers were washed with brine (10 mL), dried over Na₂SO₄, filtered and evaporated to obtain a yellow oil. The crude was purified by flash chromatography (pentane/EtOAc 1:1) to obtain the desired diol **16** (76.7 mg, 0.27 mmol, 84%, dr = 10:1) as a white solid; Mp: 126.3–126.8 °C; *R_f* = 0.25 (pentane/EtOAc 1:1); $[\alpha]_D^{25} = +45.6$ (*c* 0.16, CHCl₃); FTIR (neat) $\tilde{\nu} = 3449, 2959, 2929, 2360, 1727, 1714, 1660, 1429, 1351, 1255, 1202, 1037, 1021, 739 \text{ cm}^{-1}$; ¹H NMR (400 MHz, CDCl₃) $\delta = 5.16$ (dd, *J* = 12.2, 6.4 Hz, 1H), 5.03–4.70 (m, 2H), 3.69 (s, 1H), 3.62 (s, 3H), 2.72–2.53 (m, 1H), 2.50–2.40 (m, 1H), 2.40–2.22 (m, 3H), 2.22–2.09

(m, 1H), 1.90 (dd, $J = 9.2, 7.0$ Hz, 2H), 0.90 (d, $J = 7.1$ Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) $\delta = 173.9, 173.5, 162.6, 128.0, 76.9, 71.5, 69.1, 63.3, 51.7, 41.4, 37.5, 22.4, 19.8, 15.7$; HRMS (ESI) m/z calcd for $\text{C}_{14}\text{H}_{19}\text{O}_6$ $[\text{M} + \text{H}]^+$: 283.1176, found: 283.1178.

Methyl (3aR,5aR,6R,8R,8aS,8bS)-8,8a-dihydroxy-6-methyl-3-oxodecahydro-5aH-indeno[4,5-c]furan-5a-carboxylate (17): Diol **16** (12.5 mg, 44.3 μmol , 1.0 eq.) and Rh/C/ Al_2O_3 (5% Rh, 9.1 mg, 4.43 μmol , 0.1 eq.) were dissolved in dry EtOAc (0.3 mL) and hydrogenated at rt under a 60 bar H_2 pressure. After 3 h, the reaction was filtered over celite and vigorously washed with EtOAc. The solvent was evaporated under reduced pressure to obtain a white solid. The residue was purified by flash chromatography (pentane/EtOAc 1:1) to obtain the desired hydrogenated lactone **17** (12.6 mg, 44.3 μmol , quant.) as a white solid; Mp: 197.7–198.5 $^\circ\text{C}$; $R_f = 0.31$ (EtOAc/pentane 1:1); $[\alpha]_D^{25} = -110.8$ (c 0.12, CHCl_3); FTIR (neat) $\tilde{\nu} = 3498, 3425, 2955, 2871, 1743, 1708, 1366, 1202, 1038, 974, 716, 617$ cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) $\delta = 4.91$ (dd, $J = 9.2, 4.1$ Hz, 1H), 4.24 (dd, $J = 9.7, 8.3$ Hz, 1H), 4.00 (dd, $J = 11.5, 9.7$ Hz, 1H), 3.69 (s, 3H), 3.43 (s, 1H), 2.93–2.84 (m, 2H), 2.63–2.55 (m, 1H), 2.22 (dt, $J = 13.0, 3.6$ Hz, 1H), 2.10 (brs, 1H), 2.06 (ddd, $J = 10.6, 6.6, 3.5$ Hz, 1H), 1.93–1.75 (m, 3H), 1.56–1.47 (m, 1H), 0.77 (d, $J = 7.2$ Hz, 3H); ^{13}C NMR (126 MHz, CDCl_3) $\delta = 179.2, 174.3, 80.0, 72.5, 67.4, 59.6, 51.7, 44.1, 39.9, 39.3, 38.2, 24.9, 22.1, 14.7$; HRMS (ESI) m/z calcd for $\text{C}_{14}\text{H}_{21}\text{O}_6$ $[\text{M} + \text{H}]^+$: 285.1333, found: 285.1334; X-ray crystal structure is given in the supporting information.

Methyl (3aS,5R,7aR)-1,4-dioxo-5-((R)-4-oxobutan-2-yl)octahydroisobenzofuran-5-carboxylate (18): Diol **17** (11.3 mg, 39.7 μmol , 1.0 eq.) was dissolved in THF/water (2:1, 0.47 mL) and NaIO_4 (27.8 mg, 0.13 mmol, 3.2 eq.) was added at rt. After 1.5 h, the reaction mixture was diluted with EtOAc (2 mL) and filtered over Na_2SO_4 . The filtrate was vigorously washed with EtOAc (4 x 2 mL). The solvent was evaporated to obtain a white solid. The residue was purified by flash chromatography (pentane/EtOAc 2:1) to obtain the desired ketoaldehyde **18** (12.5 mg, 43.6 μmol , quant.) as colorless oil; $R_f = 0.38$ (EtOAc/pentane 1:1); $[\alpha]_D^{25} = -137.3$ (c 0.12, CHCl_3); FTIR (neat) $\tilde{\nu} = 2955, 1770, 1712, 1452, 1167, 998, 732, 626$ cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) $\delta = 9.71$ (s, 1H), 4.86 (dd, $J = 9.4, 1.9$ Hz, 1H), 4.19 (dd, $J = 9.4, 6.6$ Hz, 1H), 3.70 (s, 3H), 3.30–3.18 (m, 2H),

2.93–2.84 (m, 1H), 2.82–2.68 (m, 2H), 2.29–2.17 (m, 2H), 2.15–2.04 (m, 2H), 1.07 (d, J = 6.7 Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ = 205.4, 201.3, 176.7, 170.7, 66.2, 63.0, 52.5, 47.5, 45.6, 40.2, 29.1, 28.8, 18.8, 15.8.

Methyl (3aS,5R,7aR)-5-((R)-1-(1,3-dioxolan-2-yl)propan-2-yl)-1,4-dioxooctahydroisobenzofuran-5-carboxylate (19): To the diol **17** (130.0 mg, 0.45 mmol, 1.0 eq.) dissolved in a THF and water mixture (3:1, 5 mL) was added sodium periodate (332.0 mg, 1.60 mmol, 3.4 eq.). The mixture was stirred at rt for 2 h and subsequently dried by filtrating the mixture through a patch of Na_2SO_4 and washed with EtOAc. The solvent was evaporated, and the residue was obtained as colorless oil. The crude ketoaldehyde **18** was dissolved in dry benzene (4 mL), and ethylene glycol (27 μL , 0.48 mmol, 1.1 eq.) was added, followed by *p*-TSA (9.2 mg, 0.05 mmol, 0.1 eq.). The resulting solution was heated to reflux (90 °C). After 1 h 45 min, the solution was allowed to cool to rt, and was then treated with saturated NaHCO_3 solution (3 mL) followed by extraction with EtOAc three times. The combined organic layers were washed with water, dried over Na_2SO_4 and evaporated under reduced pressure. The residue was subjected to flash chromatography on silica ($\text{CH}_2\text{Cl}_2/\text{MeOH}$, 20:1) and concentrated to give protected aldehyde **19** (149.0 mg, 0.45 mmol, quant.) as a white solid; R_f = 0.63 ($\text{CH}_2\text{Cl}_2/\text{MeOH}$ 20:1); $[\alpha]_D^{25}$ = -75.5 (c 0.77, CHCl_3); FTIR (neat) $\tilde{\nu}$ = 2954, 2887, 1773, 1737, 1709, 1450, 1372, 1343, 1297, 1220, 1188, 1148, 1121, 1050, 1027, 998, 962, 734 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ = 4.95 (dd, J = 9.3, 1.8 Hz, 1H), 4.86 (dd, J = 6.4, 2.6 Hz, 1H), 4.14 (dd, J = 9.3, 6.5 Hz, 1H), 3.97–3.90 (m, 2H), 3.87–3.79 (m, 2H), 3.70 (s, 3H), 3.47 (ddd, J = 10.0, 6.4, 1.7 Hz, 1H), 3.18 (ddd, J = 10.1, 6.4, 5.1 Hz, 1H), 2.42–2.35 (m, 1H), 2.22–2.07 (m, 4H), 2.03 (ddd, J = 14.9, 9.8, 2.6 Hz, 1H), 1.54 (ddd, J = 14.9, 6.4, 0.8 Hz, 1H), 1.09 (d, J = 6.8 Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ = 205.2, 176.8, 170.9, 103.3, 66.0, 65.1, 64.9, 64.2, 52.2, 45.7, 40.9, 36.8, 30.3, 29.1, 18.6, 15.4; HRMS (ESI) m/z calcd for $\text{C}_{16}\text{H}_{23}\text{O}_7$ $[\text{M} + \text{H}]^+$: 327.1438, found: 327.1443.

Methyl (3aR,5R,7aR)-5-((R)-1-(1,3-dioxolan-2-yl)propan-2-yl)-3a-(hydroxymethyl)-1,4-dioxooctahydroisobenzofuran-5-carboxylate (20): Acetal **19** (161.0 mg, 0.49 mmol, 1.0

eq.) was dissolved in THF (2.8 mL), and formaline (37% aq. solution, 8.3 mL, 112 mmol, 226 eq.) was added, followed by aq. citrate buffer solution (8.3 mL, pH 6.0). The resulting solution was heated to 40 °C and stirred overnight. After cooling to rt, the reaction mixture was diluted with EtOAc and water. The aqueous layer was extracted three times with EtOAc, and the combined organic layers were washed with water, dried over Na₂SO₄, and evaporated under reduced pressure. The crude residue was dissolved in a CH₂Cl₂/MeOH mixture (20:1) and filtered through a pad of silica. Purification by flash chromatography (CH₂Cl₂/MeOH 20:1) gave alcohol **20** (164.3 mg, 0.46 mmol, 94%) as a white solid; *R_f* = 0.30 (CH₂Cl₂/MeOH 20:1); [α]_D²⁵ = -23.5 (*c* 0.30, CHCl₃); FTIR (neat): $\tilde{\nu}$ = 3467, 2957, 2923, 2854, 1766, 1728, 1702, 1465, 1437, 1382, 1300, 1223, 1174, 1137, 1096, 1017, 952, 881, 860, 826, 751, 713 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ = 4.87 (dd, *J* = 5.8, 3.5 Hz, 1H), 4.34 (d, *J* = 9.8 Hz, 1H), 4.13 (d, *J* = 9.8 Hz, 1H), 4.00–3.91 (m, 3H), 3.88–3.79 (m, 2H), 3.71 (s, 3H), 3.60 (d, *J* = 10.8 Hz, 1H), 3.02–2.93 (m, 1H), 2.38–2.19 (m, 3H), 2.05 (ddd, *J* = 12.9, 11.5, 2.3 Hz, 1H), 1.98–1.84 (m, 1H), 1.79–1.64 (m, 2H), 1.12 (d, *J* = 6.8 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ = 208.5, 177.5, 171.4, 103.7, 70.9, 67.0, 65.0, 64.8, 63.3, 56.6, 42.8, 35.8, 33.8, 30.5, 28.9, 20.8, 17.0; HRMS (ESI) *m/z* calcd for C₁₇H₂₅O₈ [M + H]⁺: 357.1544, found: 357.1552.

Methyl (3aR,5R,7aR)-5-((R)-1-(1,3-dioxolan-2-yl)propan-2-yl)-3a (iodomethyl)-1,4-dioxooctahydroisobenzofuran-5-carboxylate (21): Alcohol **20** (408 mg, 1.14 mmol, 1.0 eq.), triphenylphosphine (600 mg, 2.29 mmol, 2.0 eq.) and imidazole (311 mg, 4.57 mmol, 4.0 eq.) were dissolved in dry benzene (30 mL) under argon atmosphere. Iodine (580 mg, 2.29 mmol, 2.0 eq.) was added at rt and the resulting mixture was stirred for 12 hours at that temperature. The reaction was quenched with a 1:1 mixture of saturated NaHCO₃ and Na₂S₂O₃ solution and extracted three times with EtOAc. The combined organic layers were dried over Na₂SO₄, filtered and evaporated under reduced pressure. The residue was purified by flash chromatography on silica (pentane/EtOAc 5:2) giving iodide **21** (401.2 mg, 0.86 mmol, 75%) as colorless oil; *R_f* = 0.23 (pentane/EtOAc 3:1); [α]_D²⁵ = +17.9 (*c* 0.48, CHCl₃); FTIR (neat): $\tilde{\nu}$ = 2954, 2886, 2360, 1781, 1731, 1710, 1436, 1366, 1225, 1162, 1030, 947, 824, 732 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ = 4.87 (dd, *J* = 5.8, 3.7 Hz, 1H), 4.39 (d, *J* = 10.2 Hz, 1H), 4.01 (d, *J* = 10.2 Hz, 1H), 3.98–3.87

(m, 3H), 3.86–3.78 (m, 2H), 3.72 (s, 3H), 3.15 (d, $J = 10.3$ Hz, 1H), 3.14–3.08 (m, 1H), 2.49–2.38 (m, 1H), 2.33–2.24 (m, 2H), 2.08–1.98 (m, 1H), 1.86 (tdd, $J = 13.3, 8.3, 2.3$ Hz, 1H), 1.72–1.55 (m, 2H), 1.11 (d, $J = 6.8$ Hz, 3H); ^{13}C NMR (126 MHz, CDCl_3) $\delta = 203.7, 176.2, 171.1, 103.5, 72.9, 65.1, 64.8, 63.5, 55.3, 53.0, 46.9, 36.4, 35.6, 27.5, 20.8, 17.4, 11.3$; HRMS (ESI) m/z calcd for $\text{C}_{17}\text{H}_{24}\text{IO}_7$ $[\text{M} + \text{H}]^+$: 467.0561, found: 467.0557.

Methyl (3aS,5R,7aR)-5-((R)-1-(1,3-dioxolan-2-yl)propan-2-yl)-3a-methyl-1,4-dioxooctahydroisobenzofuran-5-carboxylate (22): Iodide **21** (401.0 mg, 0.86 mmol, 1.0 eq.) was dissolved in dry and degassed benzene (30 mL) under argon atmosphere. Tri-*n*-butyltin hydride (348 μL , 1.29 mmol, 1.5 eq.) and AIBN (42.0 mg, 0.26 mmol, 0.3 eq.) were added and the mixture was stirred for 3 h at 85 °C. The solvent was removed *in vacuo* and the residue was purified by flash column chromatography (pentane/EtOAc 2:1) to give the ketone **22** (242.0 mg, 0.72 mmol, 84%) as colorless oil. $R_f = 0.18$ (pentane/EtOAc 2:1); $[\alpha]_D^{25} = -68.5$ (c 0.49, MeOH); FTIR (neat) $\tilde{\nu} = 2955, 2887, 1776, 1733, 1709, 1457, 1364, 1297, 1230, 1148, 1020, 963, 823, 733$ cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) $\delta = 4.85$ (t, $J = 4.8$ Hz, 1H), 4.47 (d, $J = 9.3$ Hz, 1H), 3.95–3.84 (m, 3H), 3.83–3.75 (m, 2H), 3.64 (s, 3H), 2.61 (dd, $J = 10.3, 6.1$ Hz, 1H), 2.43–2.32 (m, 1H), 2.26–2.11 (m, 2H), 2.03–1.86 (m, 2H), 1.59–1.53 (m, 2H), 1.33 (s, 3H), 1.03 (d, $J = 6.8$ Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) $\delta = 207.7, 177.5, 171.1, 103.5, 74.6, 65.0, 64.7, 63.3, 52.6, 50.9, 46.3, 36.1, 34.8, 27.7, 24.3, 19.5, 16.8$; HRMS (ESI) m/z calcd for $\text{C}_{17}\text{H}_{25}\text{O}_7$ $[\text{M} + \text{H}]^+$: 341.1595, found: 341.1591.

Methyl (3aS,5R,7aR)-3a-methyl-1,4-dioxo-5-((R)-4-oxobutan-2-yl)octahydroisobenzofuran-5-carboxylate (23): The ketone **22** (227.0 mg, 0.67 mmol, 1.0 eq.) was dissolved in acetone (10 mL) and 1 M aq. HCl solution (5 mL) was added. The resulting mixture was stirred at rt for 17 h. The reaction was quenched by addition of a saturated aq. NaHCO_3 solution and extracted three times with EtOAc. The combined organic layers were dried over Na_2SO_4 , filtered and evaporated under reduced pressure to yield pure aldehyde **23** (195.8 mg, 0.66 mmol, 99%) as colorless oil; $R_f = 0.56$ (pentane/EtOAc 1:1); $[\alpha]_D^{25} = -87.8$ (c 0.42, CHCl_3); FTIR (neat): $\tilde{\nu} = 2956, 2360, 2339, 1776, 1717, 1457, 1385, 1295, 1234, 1155, 1022, 991, 787$ cm^{-1} ; ^1H NMR (400 MHz,

CDCl₃) δ = 9.72 (d, J = 1.7 Hz, 1H), 4.60 (d, J = 9.4 Hz, 1H), 3.90 (d, J = 9.4 Hz, 1H), 3.68 (s, 3H), 2.94 (dq, J = 9.8, 6.8, 3.0 Hz, 1H), 2.69 (t, J = 7.4 Hz, 1H), 2.67–2.60 (m, 1H), 2.52–2.43 (m, 1H), 2.31–2.19 (m, 2H), 2.08–1.94 (m, 2H), 1.36 (s, 3H), 1.00 (d, J = 6.8 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ = 208.1, 200.7, 177.3, 170.8, 74.7, 62.3, 52.8, 51.2, 46.8, 46.2, 31.3, 27.2, 24.5, 18.9, 16.6; HRMS (ESI) m/z calcd for C₁₅H₂₁O₆ [M + H]⁺: 297.1333 found: 297.1328.

*Methyl (3a*S*,5*R*,7a*R*)-5-((*R*)-4-hydroxybutan-2-yl)-3a-methyl-1,4-*

dioxooctahydroisobenzofuran-5-carboxylate (24): Aldehyde **23** (78.1 mg, 0.26 mmol, 1.0 eq.) was dissolved in hexafluoroisopropanol (1 mL). Sodium tris(hexafluoroisopropoxy)borohydride (1 M in THF, 791 μ L, 0.79 mmol, 3.0 eq.) was added at rt under nitrogen atmosphere and stirred for 15 h. The reaction was quenched by addition of saturated NH₄Cl solution and extracted three times with EtOAc. The combined organic layers were dried over Na₂SO₄, filtered and evaporated under reduced pressure. The residue was purified by flash chromatography (pentane/EtOAc 2:3) giving alcohol **24** (66.9 mg, 0.224 mmol, 85%) as colorless oil; R_f = 0.26 (EtOAc/pentane 3:2); $[\alpha]_D^{25}$ = –79.9 (c 0.18, MeOH); FTIR (neat): $\tilde{\nu}$ = 3529, 3407, 2955, 2884, 1775, 1731, 1708, 1456, 1385, 1364, 1293, 1231, 1187, 1154, 1055, 1021, 989, 916, 848, 732, 681 cm^{–1}; ¹H NMR (400 MHz, CDCl₃) δ = 4.48 (d, J = 9.3 Hz, 1H), 3.94 (d, J = 9.3 Hz, 1H), 3.74 (ddd, J = 11.0, 8.8, 5.5 Hz, 1H), 3.69 (s, 3H), 3.67–3.58 (m, 1H), 2.64 (t, J = 8.4 Hz, 1H), 2.45–2.34 (m, 1H), 2.28 (dt, J = 9.3, 4.4 Hz, 1H), 2.25–2.17 (m, 1H), 2.07–1.92 (m, 2H), 1.69–1.52 (m, 2H), 1.39–1.33 (m, 1H), 1.37 (d, J = 2.9 Hz, 3H), 0.99 (d, J = 6.8 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ = 208.1, 177.5, 171.3, 74.6, 63.2, 61.0, 52.8, 51.0, 46.5, 35.3, 34.8, 27.4, 24.4, 19.8, 16.1; HRMS (ESI) m/z calcd for C₁₅H₂₂NaO₆ [M + Na]⁺: 321.1309, found: 321.1308.

*Methyl (3a*S*,5*R*,7a*R*)-5-((*R*)-4-iodobutan-2-yl)-3a-methyl-1,4-*

dioxooctahydroisobenzofuran-5-carboxylate (25): Alcohol **24** (26.0 mg, 0.087 mmol, 1.0 eq.) was dissolved in dry benzene (5 mL) under nitrogen atmosphere. Triphenylphosphine (45.7 mg, 0.174 mmol, 2.0 eq.), imidazole (23.7 mg, 0.349 mmol,

4.0 eq.), and iodine (44.2 mg, 0.174 mmol, 2.0 eq.) were added at rt and the resulting mixture was stirred overnight. The crude mixture was filtered, washed with benzene and evaporated under reduced pressure. The resulting oil was purified by flash chromatography (pentane/EtOAc 2:1) to give iodide **25** (33.8 mg, 0.083 mmol, 95%) as a white solid; R_f = 0.44 (pentane/EtOAc 2:1); $[\alpha]_D^{25}$ = -9.8 (c 0.58, CHCl_3); FTIR (neat): $\tilde{\nu}$ = 3245, 2971, 2927, 2860, 2171, 1762, 1728, 1700, 1591, 1532, 1489, 1439, 1385, 1368, 1287, 1230, 1207, 1173, 1119, 1069, 1012, 977, 961, 922, 905, 852, 773, 764, 722, 692 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ = 4.48 (d, J = 9.3 Hz, 1H), 3.93 (d, J = 9.3 Hz, 1H), 3.71 (s, 3H), 3.33 (ddd, J = 9.8, 7.7, 4.3 Hz, 1H), 3.07 (td, J = 9.5, 7.1 Hz, 1H), 2.68–2.60 (m, 1H), 2.37–2.14 (m, 3H), 2.05–1.91 (m, 2H), 1.89–1.78 (m, 1H), 1.71–1.60 (m, 1H), 1.37 (s, 3H), 0.97 (d, J = 6.8 Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ = 207.7, 177.4, 171.0, 74.6, 63.1, 52.9, 51.0, 46.4, 39.5, 35.6, 27.6, 24.5, 19.7, 15.1, 4.4; HRMS (ESI) m/z calcd for $\text{C}_{15}\text{H}_{22}\text{IO}_5$ $[\text{M} + \text{H}]^+$: 409.0506, found: 409.0508; X-ray crystal structure is given in the supporting information.

Samarium(II) iodide: Following a literature procedure,⁴⁸ samarium (151 mg, 1.0 mmol, 1.0 eq.) was activated by hot stirring with a heat gun *in vacuo*. After cooling, 1,2-diiodoethane (prewashed by extraction with saturated aqueous $\text{Na}_2\text{S}_2\text{O}_3$ solution, 141 mg, 0.5 mmol, 1.0 eq.) was added under argon atmosphere. THF (dried over sodium, 2 mL) was added and the mixture was stirred for 5 min at room temperature. After that time, more THF (3 mL) was added and the solution was stirred until it turned dark blue and SmI_2 was obtained as a 0.1 M solution in THF and was used as such.

Methyl (3aR,5aR,6R,8aR,8bS)-8a-hydroxy-6,8b-dimethyl-3-oxodecahydro-5aH-indeno[4,5-c]furan-5a-carboxylate (26): Iodide **25** (118.0 mg, 0.289 mmol, 1.0 eq.) was dissolved in hexamethylphosphoramide (500 μL , 2.89 mmol, 10.0 eq.) under nitrogen atmosphere and cooled down to 0 °C. A freshly prepared SmI_2 solution (0.1 M in THF, 7.22 mL, 0.722 mmol, 2.5 eq.) was added and the resulting mixture was slowly heated up to rt and stirred for 1 hour. The reaction was quenched by addition of a saturated aqueous Rochelle's salt solution and stirred at that temperature for 20 min. Extraction was performed three times with EtOAc. The combined organic layers were dried over

Na₂SO₄, filtered and evaporated under reduced pressure. The residue was purified by flash chromatography on silica (pentane/EtOAc 4:1, CAM dip) giving the tricyclic product **26** (61.5 mg, 0.218 mmol, 76%) as a white solid; Mp: 97.8–98.6 °C; *R_f* = 0.41 (pentane/EtOAc 4:1); [α]_D²⁵ = –17.6 (*c* 0.29, MeOH); FTIR (neat): $\tilde{\nu}$ = 3495, 2959, 2900, 2877, 1770, 1708, 1454, 1435, 1384, 1354, 1325, 1291, 1270, 1239, 1181, 1148, 1102, 1076, 1058, 1040, 1009, 988, 966, 920, 896, 845, 768, 739, 714, 681 cm^{–1}; ¹H NMR (400 MHz, CDCl₃) δ = 5.17 (d, *J* = 2.1 Hz, 1H), 4.44 (d, *J* = 9.5 Hz, 1H), 3.97 (d, *J* = 9.5 Hz, 1H), 3.79 (s, 3H), 2.40–2.32 (m, 1H), 2.24–2.16 (m, 1H), 2.14–2.00 (m, 2H), 1.96–1.86 (m, 2H), 1.82–1.71 (m, 1H), 1.66–1.58 (m, 1H), 1.43 (td, *J* = 13.9, 2.6 Hz, 1H), 1.35–1.29 (m, 1H), 1.23 (s, 3H), 1.00 (d, *J* = 6.9 Hz, 3H); ¹³C NMR (126 MHz, CDCl₃) δ = 178.8, 178.5, 85.4, 74.6, 58.0, 52.6, 48.1, 46.8, 45.1, 36.9, 33.8, 29.4, 23.1, 22.4, 21.2; HRMS (ESI) *m/z* calcd for C₁₅H₂₂NaO₅ [*M* + Na]⁺: 305.1359, found: 305.1359.

(3*aR*,5*aR*,6*R*,8*aR*,8*bS*)-8*a*-Hydroxy-6,8*b*-dimethyl-3-oxodecahydro-5*aH*-indeno[4,5-*c*]furan-5*a*-carbaldehyde (**27**): In a dry round bottomed flask under argon atmosphere, ester **26** (19.7 mg, 0.070 mmol, 1.0 eq.) was dissolved in dry THF (2 mL) and cooled to 0 °C. Borane-methyl sulfide complex (2 M in THF, 70 μ L, 0.140 mmol, 2.0 eq.) was added, the resulting mixture was stirred for 15 min at 0 °C and subsequently at 45 °C for 5.5 h. The mixture was allowed to cool down to rt for 10 min before ethanol was added until bubbling (exothermic) ceased. The solution was evaporated giving a colorless oil. Purification was performed by flash chromatography (pentane/EtOAc 3:2) giving aldehyde **27** (11.6 mg, 0.0460 mmol, 66%) as a white solid; Mp: 51.4–51.9 °C; *R_f* = 0.32 (pentane/EtOAc 3:2, CAM dip); [α]_D²⁵ = –13.7 (*c* 0.19, MeOH); FTIR (neat) $\tilde{\nu}$ = 3484, 2957, 2876, 1748, 1704, 1460, 1373, 1287, 1227, 1158, 1140, 1003, 973, 855, 793, 732, 690, 551 cm^{–1}; ¹H NMR (400 MHz, CDCl₃) δ = 9.84 (s, 1H), 4.23 (d, *J* = 0.8 Hz, 1H), 3.95 (d, *J* = 9.6 Hz, 1H), 3.51 (s, 1H), 2.29–2.18 (m, 2H), 2.12 (dddd, *J* = 14.4, 10.4, 9.1, 2.2 Hz, 2H), 1.98–1.84 (m, 2H), 1.82–1.72 (m, 1H), 1.72–1.60 (m, 1H), 1.51–1.41 (m, 1H), 1.41–1.32 (m, 1H), 1.22 (d, *J* = 0.8 Hz, 3H), 1.08 (d, *J* = 7.0 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ = 210.1, 178.9, 86.4, 75.0, 59.8, 47.5, 46.4, 44.6, 37.8, 31.5, 30.5, 22.5, 22.3, 19.8; HRMS (ESI) *m/z* calcd for C₁₄H₂₀O₄Na [*M* + Na]⁺: 275.1254, found: 275.1253.

(3*aR*,5*aS*,6*R*,8*aR*,8*bS*,11*S*)-11-Hydroxy-6,8*b*-dimethylhexahydro-6*H*-8*a*,5*a*-(epoxyethano)indeno[4,5-*c*]furan-3,10(1*H*)-dione (**28**): In a dry round bottomed flask under argon atmosphere, ester **26** (31.8 mg, 0.113 mmol, 1.0 eq.) was dissolved in dry THF (3 mL) and cooled to 0 °C. Borane-methyl sulfide complex (2 M in THF, 113 μ L, 0.225 mmol, 2.0 eq.) was added, the resulting mixture was stirred for 15 min at 0 °C and subsequently at 45 °C for 7 h. The mixture was allowed to cool down to rt for 10 min before ethanol was added until bubbling (exothermic) ceased. The solution was evaporated giving a colorless oil. Crude aldehyde **27** was dissolved in THF/water (1:1, 4 mL) and potassium cyanide (25.8 mg, 0.396 mmol, 3.5 eq.) was added. The mixture was stirred for 2 d before EtOAc was added and the mixture was stirred for an additional day. The phases were separated and the aqueous layer was extracted with EtOAc for an additional day. Extractions were performed until no more product was observed in the separated organic layer. The combined organic layers were dried over Na₂SO₄, filtered and evaporated under reduced pressure. The residue was purified by flash chromatography (EtOAc/pentane 3:2) giving lactone **28** (20.0 mg, 0.0713 mmol, 63%) as a white solid; *R*_f = 0.33 (EtOAc/pentane 1:1); [α]_D²⁵ = -41.9 (*c* 0.16, MeOH); FTIR (neat) $\tilde{\nu}$ = 3421, 2965, 2888, 1774, 1758, 1678, 1462, 1379, 1254, 1196, 1159, 1132, 1091, 1026, 974, 944, 876, 820, 753, 716 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ = 4.40 (d, *J* = 9.6 Hz, 1H), 4.35 (d, *J* = 2.8 Hz, 1H), 3.92 (d, *J* = 9.6 Hz, 1H), 2.69 (d, *J* = 2.9 Hz, 1H), 2.47 (dt, *J* = 8.3, 4.2 Hz, 1H), 2.29–2.19 (m, 1H), 2.12–2.00 (m, 3H), 1.90–1.78 (m, 2H), 1.75–1.65 (m, 1H), 1.35 (s, 3H), 1.08 (d, *J* = 6.7 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ = 179.7, 176.8, 98.1, 75.7, 68.7, 53.5, 44.7, 44.3, 43.0, 34.5, 30.2, 25.3, 22.3, 20.2, 14.5; HRMS (ESI) *m/z* calcd for C₁₅H₂₁O₅ [*M* + *H*]⁺: 281.1384, found: 281.1380.

(3*aS*,4*R*,6*S*,6*aS*,7*R*,9*aR*,9*bS*)-7,9*b*-Dimethylhexahydro-3*H*,6*H*,7*H*-9*a*,6-(epoxymethano)-4,6*a*-methanocyclopenta[*c*]furo[3,4-*e*]oxepine-3,11-dione (**29**): Alcohol **28** (2.5 mg, 8.9 μ mol, 1.0 eq.) was dissolved in dry THF (1 mL) under nitrogen atmosphere and cooled down to -78 °C. LDA (0.28 M in THF, 96 μ L, 26.8 μ mol, 3.0 eq.) was added at -78 °C and the mixture was stirred for 1 h. A solution of phenylselenenyl bromide (4.2 mg, 17.8 μ mol, 2.0 eq.) in THF (0.15 mL) and was then added at -78 °C. The resulting mixture

was stirred for 1 h at that temperature and subsequently heated up to 0 °C and stirred for additional 30 min. The reaction mixture was diluted with a saturated NH₄Cl solution and EtOAc. The layers were separated and the aqueous layer was extracted twice with EtOAc. The combined organic layers were dried over Na₂SO₄, filtered and evaporated under reduced pressure. The residue was dissolved in THF (1 mL) and cooled down to 0 °C. One drop of AcOH and H₂O₂ (35% in water) were added and the mixture was stirred for 2 h before a saturated NaHCO₃ solution was added and stirred 1 h at rt. Extraction was performed three times with EtOAc. The combined organic layers were dried over Na₂SO₄, filtered and evaporated under reduced pressure. Purification was performed by flash chromatography to yield the desired product **29** (1.2 mg, 4.31 μmol, 48%) as a crystalline solid; *R_f* = 0.21 (pentane/EtOAc 1:4); [α]_D²⁵ = –68.4 (*c* 0.29, MeOH); FTIR (neat): $\tilde{\nu}$ = 3357, 2923, 2853, 1762, 1662, 1467, 1378, 1263, 1208, 1159, 1095, 1022, 962, 930, 902, 795, 748 cm^{–1}; ¹H NMR (400 MHz, d₄-MeOD) δ = 4.75 (t, *J* = 5.7 Hz, 1H), 4.65 (d, *J* = 9.3 Hz, 1H), 4.01 (s, 1H), 3.87 (d, *J* = 9.3 Hz, 1H), 3.07 (d, *J* = 5.2 Hz, 1H), 2.31–2.21 (m, 2H), 2.09–1.94 (m, 4H), 1.36 (s, 3H), 1.27–1.15 (m, 1H), 1.08 (d, *J* = 7.0 Hz, 3H); ¹³C NMR (101 MHz, d₄-MeOD) δ = 177.7, 174.9, 101.9, 78.7, 75.6, 75.5, 60.4, 55.4, 44.7, 39.0, 35.4, 34.0, 32.8, 27.3, 15.8; HRMS (ESI) *m/z* calcd for C₁₅H₁₉O₅ [*M* + *H*]⁺: 279.1227, found: 279.1223; X-ray crystal structure is given in the supporting information.

Methyl (1R,2R)-1-((R)-5-(benzyloxy)-4-methyl-3-oxopentyl)-2-methyl-5-oxocyclopentane-1-carboxylate (35): To a solution of β -keto ester **7** (10 mg, 0.064 mmol, 1.0 eq) and enone **34** (19.6 mg, 0.096 mmol, 1.2 eq) in acetone (0.25 mL) was added K₂CO₃ (4.4 mg, 0.032 mmol, 0.5 eq) and the mixture was heated to 40 °C overnight. After cooling down to room temperature the mixture was filtered through a short pad of celite. The solvent was removed and the residue was subjected to flash column chromatography (pentane/Et₂O 5:1 to 3:1) to give ester **35** (16.2 mg, 45 μmol, 70%) as colorless oil; *R_f* = 0.53 (pentane/Et₂O 6:1); [α]_D²⁴ = +27.3 (*c* 0.44, CHCl₃); FTIR (neat): $\tilde{\nu}$ = 2960, 2877, 2361, 1729, 1496, 1455, 1375, 1236, 1168, 1101, 1026, 997, 741, 699 cm^{–1}; ¹H NMR (400 MHz, CDCl₃) δ = 7.36–7.26 (m, 5H), 4.48 (d, *J* = 3.2 Hz, 2H), 3.68 (s, 3H), 3.62

(dd, $J = 9.1, 7.7$ Hz, 1H), 3.49–3.44 (m, 1H), 2.91–2.78 (m, 2H), 2.59–2.46 (m, 2H), 2.24–2.08 (m, 3H), 2.06–1.87 (m, 2H), 1.79–1.66 (m, 1H), 1.07 (d, $J = 7.0$ Hz, 3H), 1.01 (d, $J = 6.8$ Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) $\delta = 216.5, 212.4, 171.0, 138.3, 128.5, 127.7, 127.7, 73.4, 72.5, 62.0, 51.9, 46.7, 41.6, 38.7, 37.2, 28.3, 26.1, 16.0, 13.7$; HRMS (ESI) m/z calcd for $\text{C}_{21}\text{H}_{28}\text{NaO}_5$ $[\text{M} + \text{Na}]^+$: 383.1829, found: 383.1834.

Methyl (1R,2R)-1-((R)-5-hydroxy-4-methyl-3-oxopentyl)-2-methyl-5-oxocyclopentane-1-carboxylate (36): A mixture of ester **35** (50.0 mg, 0.139 mmol, 1.0 eq) and 5 % Pd/C (44.3 mg, 0.021 mmol, 15 mol%) in EtOH (3.0 mL) was stirred under a hydrogen atmosphere. After completion of the reaction (2.5 h), the mixture was filtered through a short pad of celite and the solvent was removed under reduced pressure to give alcohol **36** (37.5 mg, 0.139 mmol, quant.) as colorless oil; $R_f = 0.16$ (pentane/Et₂O 1:2); $[\alpha]_D^{26} = +40.0$ (c 0.98, CHCl_3); FTIR (neat): $\tilde{\nu} = 3732, 3505, 2960, 2881, 2361, 2340, 1728, 1459, 1381, 1237, 1169, 1124, 1035, 678$ cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) $\delta = 3.69$ (s, 3H), 3.75–3.63 (m, 2H), 2.87 (ddd, $J = 18.1, 9.2, 5.7$ Hz, 1H), 2.80–2.70 (m, 1H), 2.64–2.47 (m, 2H), 2.33–2.03 (m, 5H), 1.91 (ddd, $J = 14.7, 9.3, 5.7$ Hz, 1H), 1.80–1.66 (m, 1H), 1.10 (d, $J = 7.2$ Hz, 3H), 1.02 (d, $J = 6.8$ Hz, 3H); ^{13}C NMR (63 MHz, CDCl_3) $\delta = 216.8, 214.1, 170.9, 64.7, 61.9, 52.0, 48.3, 42.4, 38.7, 37.0, 28.3, 26.5, 16.0, 13.3$; HRMS (ESI) m/z calcd for $\text{C}_{14}\text{H}_{22}\text{NaO}_5$ $[\text{M} + \text{Na}]^+$: 293.1359, found: 293.1361.

Methyl (1R,2R)-2-methyl-1-((E)-4-methyl-5-oxopent-3-en-1-yl)-5-oxocyclopentane-1-carboxylate (38): To a solution of β -keto ester **7** (284 mg, 1.82 mmol, 1.0 eq) and $\alpha,\beta,\gamma,\delta$ -unsaturated aldehyde **37** (4.0 mL, 2.00 mmol, 1.1 eq, 0.5 M solution in CH_2Cl_2) in acetone (10.0 mL) was added K_2CO_3 (503 mg, 3.64 mmol, 2.0 eq) and the mixture was heated to 30 °C overnight. After cooling to room temperature, the mixture was filtered through a short pad of celite. The solvent was removed and the residue was subjected to flash column chromatography (pentane/Et₂O 3:1 to 2:1) to give **38** (319 mg, 1.26 mmol, 70%) as colorless oil; $R_f = 0.47$ (pentane/Et₂O 1:1); $[\alpha]_D^{24} = +55.2$ (c 1.01, CHCl_3); FTIR (neat): $\tilde{\nu} = 2958, 2361, 1730, 1685, 1645, 1436, 1406, 1356, 1232, 1168, 1124, 1042, 994, 669$ cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) $\delta = 9.36$ (s, 1H), 6.43 (td, $J = 7.3, 1.4$ Hz, 1H), 3.69 (s, 3H), 2.63–2.51 (m, 2H), 2.30–2.18 (m, 3H), 2.15–1.96 (m, 2H), 1.84–1.74

(m, 2H), 1.72 (d, $J = 1.1$ Hz, 3H), 1.05 (d, $J = 6.9$ Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) $\delta = 216.0, 195.1, 170.6, 153.4, 139.7, 62.4, 51.9, 40.6, 38.6, 30.7, 28.1, 23.9, 15.8, 9.2$; HRMS (ESI) m/z calcd for $\text{C}_{14}\text{H}_{20}\text{NaO}_4$ $[\text{M} + \text{Na}]^+$: 275.1254, found: 275.1255.

Methyl (1R,5S,7R)-7-methyl-8-oxo-4-(1-oxopropan-2-yl)bicyclo[3.2.1]octane-1-

carboxylate (39): A 0.2 M stock solution of $\text{Me}_3\text{AlSPhLi}$ was prepared by adding Me_3Al (2.0 M in toluene, 0.5 mL, 1.0 mmol) to a solution of PhSH (111 mg, 1.0 mmol) and *n*-BuLi (1.6 M in hexane, 0.63 mL, 1.0 mmol) in THF (3.8 mL) at 0 °C. To a solution of α,β -unsaturated aldehyde **38** (50 mg, 0.198 mmol, 1.0 eq) in THF was added the stock solution (1.28 mL, 0.257 mmol, 1.3 eq) and the mixture was stirred at 40 °C for 3 h before it was poured into saturated NH_4Cl solution. The layers were separated and the aqueous was extracted with CH_2Cl_2 (3 x). The combined organic layers were washed with water, dried over Na_2SO_4 , filtered and concentrated. The residue was subjected to flash column chromatography (pentane/ Et_2O 1:1) to give **39** as colorless oil (18.4 mg, 0.073 mmol, 37%, dr = 2:1); $R_f = 0.28$ (pentane/ Et_2O 1:1); FTIR (neat): $\tilde{\nu} = 3487, 2954, 2875, 1750, 1725, 1450, 1380, 1341, 1295, 1274, 1231, 1194, 1030, 692\text{ cm}^{-1}$; ^1H NMR (400 MHz, CDCl_3) $\delta = 9.54$ (d, $J = 3.3$ Hz, 1H), 9.49 (d, $J = 3.3$ Hz, 1H), 3.72 (s, 6H), 2.50–2.08 (m, 12H), 1.98–1.87 (m, 2H), 1.82–1.70 (m, 2H), 1.62–1.48 (m, 2H), 1.44–1.35 (m, 2H), 1.09 (d, $J = 6.6$ Hz, 3H), 1.05 (d, $J = 6.8$ Hz, 3H), 0.99 (d, $J = 7.1$ Hz, 3H), 0.97 (d, $J = 7.1$ Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) $\delta = 213.9, 213.4, 203.4, 203.1, 171.0, 170.9, 60.9, 60.4, 51.9, 49.8, 49.5, 49.1, 48.1, 46.7, 46.5, 38.2, 38.0, 36.2, 36.1, 28.8, 28.5, 23.5, 21.7, 21.1, 21.1, 12.3, 12.2$; HRMS (ESI) m/z calcd for $\text{C}_{14}\text{H}_{20}\text{NaO}_4$ $[\text{M} + \text{Na}]^+$: 275.12538, found: 275.12529.

Methyl (1R,2R)-2-methyl-5-oxo-1-(3-oxopropyl)cyclopentane-1-carboxylate (40): To a solution of β -keto ester **7** (95 mg, 0.608 mmol, 1.0 eq) in CH_2Cl_2 (5.0 mL) was added NEt_3 (128 μL , 0.912 mmol, 1.5 eq), followed by acrolein (41 μL , 0.730 mmol, 1.2 eq). The mixture was stirred at room temperature for 4 h. The clear solution was filtered through a short pad of silica gel and washed with Et_2O . The solvent was evaporated and the residue subjected to flash column chromatography (pentane/ Et_2O 2:1) to give **40** (92.0 mg, 0.433 mmol, 71%) as colorless oil; $R_f = 0.36$ (pentane/ Et_2O 1:1); $[\alpha]_D^{26} = +64.7$

(*c* 1.04, CHCl₃); FTIR (neat): $\tilde{\nu}$ = 2958, 2839, 2727, 2361, 1722, 1437, 1386, 1236, 1166, 1126, 1062, 996, 836, 763, 668 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ = 9.74 (s, 1H), 3.68 (s, 3H), 2.91–2.76 (m, 1H), 2.60–2.45 (m, 2H), 2.31–2.00 (m, 4H), 1.99–1.84 (m, 1H), 1.83–1.67 (m, 1H), 1.03 (d, *J* = 6.7 Hz, 1H); ¹³C NMR (101 MHz, CDCl₃) δ = 216.3, 201.5, 170.7, 61.7, 52.1, 41.9, 39.3, 38.6, 28.3, 24.8, 15.9; HRMS (ESI) *m/z* calcd for C₁₁H₁₆NaO₄ [M + Na]⁺: 235.0941, found: 235.0940.

Methyl (1R,2R)-1-(3-hydroxy-4-methylpent-4-en-1-yl)-2-methyl-5-oxocyclopentane-1-carboxylate (41): To a solution of aldehyde **40** (61.0 mg, 0.287 mmol, 1.0 eq) in THF (0.8 mL) at –78°C was added isopropylmagnesium bromide solution (0.5 M in THF, 0.69 mL, 0.345 mmol, 1.2 eq). The mixture was allowed to stir at –78°C for 3 h, and was then quenched by addition of sat. aq. NH₄Cl solution. Water and Et₂O were added after warming up to room temperature and the layers were separated. The aqueous layer was extracted with Et₂O (2 x) and the combined organic layers were washed with brine, dried over Na₂SO₄, filtered and concentrated. The residue was subjected to flash column chromatography (pentane/Et₂O 1:1) to give **41** (24.0 mg, 0.094 mmol, 33%) as colorless oil; *R_f* = 0.57 (pentane/Et₂O 1:2); [α]_D²⁵ = +54.2 (*c* 0.78, CHCl₃); FTIR (neat): $\tilde{\nu}$ = 3481, 2958, 2879, 2361, 1730, 1451, 1382, 1329, 1234, 1165, 1125, 1061, 996, 900, 678 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ = 4.97–4.93 (m, 1H), 4.87–4.82 (m, 1H), 4.02 (t, *J* = 6.3 Hz, 1H), 3.67 (s, 3H), 2.60–2.49 (m, 1H), 2.31–2.12 (m, 2H), 2.12–2.00 (m, 1H), 1.85–1.62 (m, 5H), 1.70 (s, 3H), 1.48–1.35 (m, 1H), 1.04 (d, *J* = 6.8 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ = 216.5, 171.1, 147.0, 111.5, 76.0, 62.7, 51.8, 40.1, 38.8, 29.3, 28.2, 27.7, 17.5, 15.8; HRMS (ESI) *m/z* calcd for C₁₄H₂₂NaO₄ [M + Na]⁺: 277.1410, found: 277.1412.

Methyl (1R,2R)-1-(3-hydroxypent-4-en-1-yl)-2-methyl-5-oxocyclopentane-1-carboxylate (42): To a solution of aldehyde **40** (50.0 mg, 0.236 mmol, 1.0 eq) in THF (1.0 mL) at –78°C was added vinylmagnesium bromide solution (0.7 M in THF, 0.42 mL, 0.294 mmol, 1.25 eq). The mixture was allowed to stir at –78°C for 4 h, and was then quenched by addition of sat. aq. NH₄Cl solution. Water and Et₂O were added after warming up to room temperature and the layers were separated. The aqueous layer was

extracted with Et₂O (2 x) and the combined organic layers were washed with brine, dried over Na₂SO₄, filtered and concentrated. The residue was subjected to flash column chromatography (pentane/Et₂O 1:1) to give **42** (12.0 mg, 0.050 mmol, 21%) as colorless oil; *R_f* = 0.44 (pentane/Et₂O 1:3); [α]_D²⁵ = +62.6 (*c* 0.52, CHCl₃); FTIR (neat): $\tilde{\nu}$ = 3443, 3081, 2957, 2879, 2361, 1729, 1435, 1404, 1328, 1234, 1165, 1121, 1061, 994, 924, 883, 669 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ = 5.84 (ddd, *J* = 17.1, 10.4, 6.0 Hz, 1H), 5.23 (dt, *J* = 17.2, 1.4 Hz, 1H), 5.11 (dt, *J* = 10.4, 1.3 Hz, 1H), 4.06 (q, *J* = 5.7 Hz, 1H), 3.67 (s, 3H), 2.54 (ddd, *J* = 18.9, 8.6, 1.1 Hz, 1H), 2.30–2.13 (m, 2H), 2.11–2.01 (m, 1H), 1.92–1.84 (m, 1H), 1.83–1.61 (m, 4H), 1.44–1.34 (m, 1H), 1.04 (d, *J* = 6.9 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ = 216.6, 171.3, 140.8, 115.1, 73.2, 62.9, 51.9, 40.2, 38.9, 31.6, 28.3, 27.7, 15.9; HRMS (ESI) *m/z* calcd for C₁₃H₂₀NaO₄ [M + Na]⁺: 263.1254, found: 263.1255.

Methyl (1R,2R)-1-(3-((4-methoxybenzyl)oxy)pent-4-en-1-yl)-2-methyl-5-oxocyclopentane-1-carboxylate (43): To a solution of **42** (10.4 mg, 0.043 mmol, 1.0 eq) and 4-Methoxybenzyl-2,2,2-trichloroacetimidate (24.5 mg, 0.087 mmol, 2.0 eq) in CH₂Cl₂ (1.0 mL) at 0 °C was added Sc(OTf)₃ (1.1 mg, 4 μ mol, 5 mol%). The mixture was stirred for 1.5 h at 0 °C and then quenched with sat. aq. NaHCO₃ solution and diluted with Et₂O. The layers were separated and the aqueous was extracted with Et₂O (3 x). The combined organic layers were dried over Na₂SO₄, filtered and concentrated. The residue was subjected to flash column chromatography (pentane/Et₂O 2:1) to give **43** (13.7 mg, 0.038 mmol, 88%) as colorless oil; *R_f* = 0.53 (pentane/Et₂O 1:1); [α]_D²⁶ = +28.7 (*c* 0.36, CHCl₃); FTIR (neat): $\tilde{\nu}$ = 3728, 3728, 3703, 3629, 3600, 2956, 2361, 2340, 1749, 1613, 1513, 1459, 1301, 1246, 1174, 1065, 1035, 995, 823, 679 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ = 7.26–7.22 (m, 2H), 6.86 (dd, *J* = 8.8, 2.4 Hz, 2H), 5.69 (ddd, *J* = 17.1, 10.6, 7.6 Hz, 1H), 5.24 (d, *J* = 2.1 Hz, 1H), 5.20 (d, *J* = 8.7 Hz, 1H), 4.50 (d, *J* = 11.3 Hz, 1H), 4.28 (d, *J* = 11.3 Hz, 1H), 3.80 (s, 3H), 3.72–3.67 (m, 1H), 3.66 (s, 3H), 2.52 (dd, *J* = 18.7, 8.2 Hz, 1H), 2.32–2.21 (m, 1H), 2.21–2.11 (m, 1H), 2.08–1.98 (m, 1H), 1.90–1.66 (m, 4H), 1.36–1.26 (m, 1H), 1.02 (d, *J* = 6.8 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ = 216.5, 171.3, 159.2, 138.8, 130.9, 129.5, 117.7, 113.9, 80.6, 69.9, 63.0, 55.4, 51.9, 39.6,

39.1, 30.2, 28.3, 27.6, 16.0; HRMS (ESI) m/z calcd for $C_{21}H_{28}NaO_5$ $[M + Na]^+$: 383.1829, found: 383.1831.

Methyl (1R,2R)-2-methyl-1-(4-methyl-3-oxopent-4-en-1-yl)-5-oxocyclopentane-1-carboxylate (44): Dess-Martin periodinane (25 mg, 0.059 mmol, 1.5 eq) was added to a cold (0 °C) solution of allylic alcohol **41** (10.0 mg, 0.039 mmol, 1.0 eq) in CH_2Cl_2 (1.0 mL). The mixture was stirred at room temperature for 1.5 h and then quenched by addition of sat. aq. $Na_2S_2O_3$ and sat. aq. $NaHCO_3$ solution. The layers were separated and the aqueous layer was extracted with CH_2Cl_2 (3 x). The combined organic layers were washed with water and brine, dried over Na_2SO_4 , filtered and concentrated. The residue was subjected to flash column chromatography (pentane/ Et_2O 2:1) to give enone **44** (8.0 mg, 0.032 mmol, 81%) as colorless oil; R_f = 0.86 (pentane/ Et_2O 1:2); $[\alpha]_D^{25}$ = +31.7 (c 0.40, $CHCl_3$); FTIR (neat): $\tilde{\nu}$ = 2958, 2362, 2340, 1730, 1677, 1631, 1455, 1375, 1337, 1236, 1168, 1123, 1094, 1068, 1037, 939 cm^{-1} ; 1H NMR (400 MHz, $CDCl_3$) δ = 5.99 (s, 1H), 5.76 (d, J = 1.6 Hz, 1H), 3.69 (s, 3H), 3.06 (ddd, J = 16.9, 10.8, 4.9 Hz, 1H), 2.74 (ddd, J = 16.9, 10.9, 5.0 Hz, 1H), 2.54 (ddd, J = 18.9, 8.5, 1.9 Hz, 1H), 2.32–2.13 (m, 3H), 2.14–2.00 (m, 1H), 1.92 (ddd, J = 14.4, 10.8, 4.9 Hz, 1H), 1.86 (s, 3H), 1.83–1.67 (m, 1H), 1.04 (d, J = 6.8 Hz, 3H); ^{13}C NMR (101 MHz, $CDCl_3$) δ = 216.6, 171.3, 147.1, 111.7, 76.1, 62.8, 51.9, 40.2, 38.9, 29.5, 28.3, 27.9, 17.6, 16.0; HRMS (ESI) m/z calcd for $C_{14}H_{20}NaO_4$ $[M + Na]^+$: 275.12538, found: 275.12535.

Methyl (1R,2R)-2-methyl-5-oxo-1-(3-oxopent-4-en-1-yl)cyclopentane-1-carboxylate (45): Dess-Martin periodinane (38.2 mg, 90 μ mol, 1.5 eq) was added to a cold (0 °C) solution of allylic alcohol **42** (14.4 mg, 0.06 mmol, 1.0 eq) in CH_2Cl_2 (1.0 mL). The mixture was stirred at room temperature for 4 h and then quenched by addition of sat. aq. $Na_2S_2O_3$ and sat. aq. $NaHCO_3$ solution. The layers were separated and the aqueous was extracted with CH_2Cl_2 (3 x). The combined organic layers were washed with water and brine, dried over Na_2SO_4 , filtered and concentrated. The residue was subjected to flash column chromatography (pentane/ Et_2O 2:1) to give enone **45** (9.4 mg, 39 μ mol, 65%) as colorless oil; R_f = 0.53 (pentane/ Et_2O 1:1); $[\alpha]_D^{25}$ = +39.6 (c 0.47, $CHCl_3$); FTIR (neat): $\tilde{\nu}$ = 2959, 2361, 1728, 1617, 1436, 1404, 1236, 1168, 1123, 1067, 992, 907 cm^{-1} ; 1H NMR (400

MHz, CDCl₃) δ = 6.34 (dd, J = 17.7, 10.2 Hz, 1H), 6.24 (dd, J = 17.6, 1.5 Hz, 1H), 5.83 (dd, J = 10.2, 1.5 Hz, 1H), 3.70 (s, 3H), 3.00 (ddd, J = 17.3, 10.7, 4.9 Hz, 1H), 2.66 (ddd, J = 17.3, 10.7, 5.0 Hz, 1H), 2.54 (ddd, J = 19.0, 8.6, 1.8 Hz, 1H), 2.32–2.14 (m, 3H), 2.13–2.03 (m, 1H), 1.92 (ddd, J = 14.5, 10.8, 4.9 Hz, 1H), 1.84–1.68 (m, 1H), 1.05 (d, J = 6.8 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ = 216.5, 200.2, 170.9, 136.5, 128.4, 61.9, 52.0, 42.2, 38.7, 34.7, 28.4, 26.9, 16.0; HRMS (ESI) m/z calcd for C₁₃H₁₈NaO₄ [M + Na]⁺: 261.1097, found: 261.1099.

Methyl (1R,2R)-1-((E)-3-hydroxy-6-(trimethylsilyl)hex-4-en-1-yl)-2-methyl-5-oxocyclopentane-1-carboxylate (46): To a degassed solution of allylic alcohol **42** (27.9 mg, 0.120 mmol, 1.0 eq) and allyltrimethyl silane (137 mg, 0.120 mmol, 10 eq) in CH₂Cl₂ (1.5 mL) was added Grubbs 2nd generation catalyst (5.00 mg, 6 μ mol, 5.0 mol%). The mixture was heated for 4 h at 40 °C and then filtered through a short pad of silica (eluted with CH₂Cl₂). The solvent was removed under reduced pressure and the residue was subjected to flash column chromatography (pentane/Et₂O 2:1 to 1:1) to give **46** (20.1 mg, 0.062 mmol, 51%) as colorless oil; R_f = 0.54 (pentane/Et₂O 1:3); $[\alpha]_D^{26}$ = +30.5 (c 1.01, CHCl₃); FTIR (neat): $\tilde{\nu}$ = 3444, 2955, 2361, 1731, 1457, 1405, 1245, 1162, 1122, 1059, 996, 966, 849, 755, 695 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ = 5.63 (dt, J = 15.6, 8.2 Hz, 1H), 5.28 (dd, J = 15.2, 7.4 Hz, 1H), 3.98 (q, J = 6.7 Hz, 1H), 3.66 (s, 3H), 2.53 (dd, J = 18.8, 8.5 Hz, 1H), 2.31–2.13 (m, 2H), 2.09–2.01 (m, 1H), 1.88–1.61 (m, 5H), 1.45 (d, J = 8.2 Hz, 2H), 1.35–1.26 (m, 1H), 1.03 (d, J = 6.8 Hz, 3H), -0.01 (s, 9H); ¹³C NMR (101 MHz, CDCl₃) δ = 216.5, 171.3, 131.2, 129.6, 73.7, 62.9, 51.9, 40.0, 39.0, 32.0, 28.3, 27.9, 22.9, 15.9, -1.8; HRMS (ESI) m/z calcd for C₁₇H₃₀NaO₄Si [M + Na]⁺: 349.1806, found: 349.1805.

Methyl (1R,2R)-2-methyl-5-oxo-1-((E)-3-oxo-6-(trimethylsilyl)hex-4-en-1-yl)cyclopentane-1-carboxylate (48): To a degassed solution of enone **45** (9.4 mg, 0.039 mmol, 1.0 eq) and allyltrimethyl silane (45 mg, 0.39 mmol, 10 eq) in CH₂Cl₂ (0.5 mL) was added Grubbs 2nd generation catalyst (1.65 mg, 2 μ mol, 5.0 mol%). The mixture was heated for 1 h at 40 °C and then filtered through a short pad of silica (eluted with CH₂Cl₂). The solvent was removed under reduced pressure and the residue was

subjected to flash column chromatography (pentane/Et₂O 3:1) to give **48** (9.2 mg, 0.028 mmol, 73%) as colorless oil; R_f = 0.58 (pentane/Et₂O 1:1); $[\alpha]_D^{25}$ = +21.6 (*c* 0.46, CHCl₃); FTIR (neat): $\tilde{\nu}$ = 2956, 2921, 2361, 1730, 1660, 1613, 1436, 1408, 1380, 1328, 1292, 1246, 1196, 1167, 1125, 1040, 979, 847, 756, 698, 669 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ = 6.97 (dt, *J* = 15.5, 8.8 Hz, 1H), 5.94 (dt, *J* = 15.7, 1.2 Hz, 1H), 3.69 (s, 3H), 2.84 (ddd, *J* = 16.1, 10.9, 4.9 Hz, 1H), 2.59–2.49 (m, 2H), 2.30–2.02 (m, 4H), 1.98–1.88 (m, 1H), 1.80–1.71 (m, 3H), 1.04 (d, *J* = 6.8 Hz, 3H), 0.05 (s, 9H); ¹³C NMR (101 MHz, CDCl₃) δ = 216.5, 199.3, 171.0, 147.0, 128.6, 62.1, 52.0, 41.8, 38.7, 35.0, 28.3, 27.3, 25.4, 16.0, -1.6; HRMS (ESI) *m/z* calcd for C₁₇H₂₈NaO₄Si [M + Na]⁺: 347.1649, found: 347.1650.

Methyl (1R,2R)-2-methyl-1-(2-(4-methyl-2-oxo-2,5-dihydrofuran-3-yl)ethyl)-5-oxocyclopentane-1-carboxylate (49): Ester **11** (516.0 mg, 1.25 mmol, 1.0 eq.) and Fe(acac)₃ (22.0 mg, 62.3 μ mol, 5 mol%) were added to a dry and degassed solution of THF (12.0 mL) and NMP (0.6 mL) and the reaction mixture was cooled to -30 °C. The mixture was stirred for 10 minutes at this temperature and MeMgBr (3.2 M in THF, 545 μ l, 1.74 mmol, 1.4 eq.) was rapidly added to the orange solution. The color of the mixture turned into a brown to green mixture, which finally resulted in a colorless solution with a brown precipitate. The reaction was quenched after 1 h with saturated NH₄Cl solution at -30 °C. The mixture was allowed to warm up to rt, water was added and the organic layer was separated. The aqueous layer was extracted with EtOAc (4 x 20 mL) and the combined organic layers were dried over Na₂SO₄, filtered and evaporated under reduced pressure. The residue was purified by flash chromatography (pentane/EtOAc 2:1 then 1:1) to obtain the desired β -methyl lactone **49** (333.9 mg, 1.19 mmol, 96%) as colorless oil; R_f = 0.24 (EtOAc/pentane 1:2); $[\alpha]_D^{25}$ = +61.0 (*c* 0.29, CHCl₃); FTIR (neat): $\tilde{\nu}$ = 2985, 1743, 1678, 1451, 1228, 1166, 1032, 762 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ = 4.61 (s, 2H), 3.69 (s, 3H), 2.59–2.45 (m, 2H), 2.42–2.32 (m, 1H), 2.29–2.08 (m, 3H), 2.05 (brs, 3H), 1.92–1.87 (m, 2H), 1.85–1.73 (m, 1H), 1.07 (d, *J* = 6.8 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ = 216.4, 174.9, 171.0, 157.7, 127.0, 72.7, 62.7, 52.0, 40.3, 38.9, 30.6, 28.4, 18.4, 16.1, 12.3; HRMS (ESI) *m/z* calcd for C₁₅H₂₁O₅ [M + H]⁺: 281.1384, found: 281.1383.

Methyl (3aR,5aR,6R,8aR,8bS)-8a-hydroxy-6,8b-dimethyl-3-oxodecahydro-5aH-indeno[4,5-c]furan-5a-carboxylate (26) and *methyl (1R,5R)-2-hydroxy-5-methyl-1-(2-(4-methyl-2-oxo-2,5-dihydrofuran-3-yl)ethyl)cyclopentane-1-carboxylate (66)*: The methylester **49** (40.4 mg, 144 mmol, 1.0 eq) was dissolved in dry THF (0.59 ml) and water (17.6 ml) (both solvents were separately freeze-thawed, three cycles). SmI₂ (3.60 ml, 360 mmol, 2.5 eq, 0.1 M solution in THF, purchased from Sigma-Aldrich) was added over 1 h at room temperature by syringe-pump and stirred for 12 h at room temperature. The colorless solution was quenched with Rochelle's salt (saturated, 1.3 ml) and stirred for 30 minutes at rt. The mixture was extracted with EtOAc (4 x 10 ml) and the combined organic layers were washed with brine (1 x 5 ml), dried over MgSO₄, filtered and evaporated to obtain a yellow oil. The mixture was purified by flash column chromatography (5:1, 4:1 then 2:1) to obtain the desired cyclized product **26** as white solid (20.5 mg, 50.4 mmol, 50%, dr 5:2) and reduced alcohol **66** (see supporting information) as colorless oil (15.8 mg, 38.9 mmol, 39%, dr = 10:1); the analytical data for the tricyclic product **26** matched those reported above. Analytical data for the alcohol byproduct **66**: colorless oil (39%); *R_f* = 0.17 (pentane/EtOAc 2:1); [α]_D²⁶ = -47.4 (*c* 1.53, CHCl₃); FTIR (neat): $\tilde{\nu}$ = 3468, 2956, 2869, 1721, 1676, 1448, 1261, 1191, 1091, 1036, 802 cm⁻¹; ¹H NMR (600 MHz, CDCl₃) δ = 4.63 (bs, 2H), 4.58 (dd, *J* = 6.6, 2.9 Hz, 1H), 3.70 (s, 3H), 2.38–2.31 (m, 2H), 2.19 (t, *J* = 11.8 Hz, 1H) 2.09–2.04 (m, 1H), 2.04 (s, 3H), 1.99–1.92 (m, 2H), 1.66–1.60 (m, 2H), 1.25–1.18 (m, 1H), 0.89 (d, *J* = 6.9 Hz, 3H); ¹³C NMR (150 MHz, CDCl₃) δ = 176.1, 174.7, 156.6, 127.4, 75.3, 73.0, 61.5, 51.3, 41.6, 32.7, 30.8, 29.7, 19.8, 15.8, 12.3; HRMS (ESI) *m/z* calcd for C₁₅H₂₂O₅Na [M + Na]⁺: 305.1359, found: 305.1359.

Methyl (1R,2R)-1-(2-((tert-butyldimethylsilyl)oxy)ethyl)-2-methyl-5-(propan-2-ylidene)-cyclopentane-1-carboxylate (51): Freshly distilled diisopropylamine (20.0 mL, 135 mmol, 1.4 eq.) was added to dry THF (50 mL) and cooled to -78 °C. *n*-BuLi (1.6 M in hexane, 79.0 mL, 126 mmol, 1.3 eq.) was slowly added to the solution and stirred for 15 min at this temperature. The reaction mixture was allowed to warm up to 0 °C and was stirred for 1 h at this temperature. The solution was cooled to -78 °C, methyl (2R)-2-

methyl-5-(propan-2-ylidene)cyclopentane-1-carboxylate (17.6 g, 97.0 mmol, 1.0 eq.) and dry DMPU (62.0 g, 59.0 mL, 483 mmol, 5.0 eq.) were added slowly to the solution and stirred for 15 min at this temperature and then for 1 h at 0 °C. The yellow solution was cooled to –78 °C and a solution of tert-butyl(2-iodoethoxy)dimethylsilane⁴¹ (33.2 g, 116 mmol, 1.2 eq.) in dry THF (50 mL) was added to the reaction mixture. The mixture was stirred for 1 h at this temperature and then was allowed to warm up to rt over 1 h. The reaction was quenched with aq. HCl soln. (1 M, 200 mL). The aqueous layer was extracted with Et₂O (4 x 150 mL). The combined organic layers were washed with brine (100 mL), dried over Na₂SO₄, filtered and the solvent was removed under reduced pressure. The residue was purified by flash column chromatography (pentane/Et₂O 40:1, then 20:1 and 8:1) to afford the alkylated ester **51** (25.3 g, 74.4 mmol, 77%) as yellow oil; R_f = 0.24 (pentane/Et₂O 40:1); $[\alpha]_D^{26}$ = +37.8 (*c* 0.79, CHCl₃); FTIR (neat): $\tilde{\nu}$ = 2953, 2933, 2858, 1727, 1462, 1434, 1253, 1218, 1186, 1155, 1095, 1004, 837, 775, 631 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ = 3.63 (s, 3H), 3.62–3.53 (m, 2H), 2.49 (dd, *J* = 15.4, 7.3 Hz, 1H), 2.29–2.17 (m, 2H), 2.17–2.04 (m, 1H), 1.96 (ddd, *J* = 14.2, 9.0, 6.4 Hz, 1H), 1.77–1.69 (m, 1H), 1.64 (d, *J* = 1.7 Hz, 3H), 1.62–1.53 (m, 1H), 1.50 (d, *J* = 1.7 Hz, 3H), 0.90–0.89 (m, 3H), 0.88 (s, 9H), 0.03 (d, *J* = 1.7 Hz, 6H); ¹³C NMR (101 MHz, CDCl₃) δ = 176.6, 137.7, 125.0, 60.4, 57.2, 51.4, 44.6, 36.9, 32.6, 32.2, 26.1, 22.4, 20.6, 18.4, 14.9, –5.1, –5.1; HRMS (ESI) *m/z* calcd for C₁₉H₃₇O₃Si [M + H]⁺ 341.2506, found: 341.2507.

Methyl (1R,2R)-1-(2-hydroxyethyl)-2-methyl-5-(propan-2-ylidene)cyclopentane-1-carboxylate (52): The protected alcohol **51** (2.32 g, 6.81 mmol, 1.0 eq.) was dissolved in dry THF (50 mL) and cooled to 0 °C. Acetic acid (1.0 mL) and TBAF (1.0 M in THF, 6.81 mL, 6.81 mmol, 1.0 eq.) were slowly added. The solution was allowed to warm up to rt and stirred for 18 h. The solution was cooled to 0 °C and quenched with aq. sat. NaHCO₃ solution (80 mL). The aqueous layer was extracted with Et₂O (3 x 100 mL) and the combined organic layers were washed with brine (50 mL). The organic layer was dried over Na₂SO₄, filtered and evaporated under reduced pressure. The crude was purified by flash column chromatography (pentane/Et₂O 1:1) to obtain the alcohol **52** (1.22 g, 5.39 mmol, 79%) as a slightly yellow oil; R_f = 0.27 (pentane/Et₂O 1:1); $[\alpha]_D^{26}$ = +35.4 (*c* 0.43, CHCl₃); FTIR (neat): $\tilde{\nu}$ = 3428, 2952, 2334, 1725, 1457, 1226, 1032, 891,

768 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ = 3.75–3.62 (m, 2H), 3.66 (s, 3H), 2.52 (dd, J = 15.6, 7.1 Hz, 1H), 2.29–1.99 (m, 5H), 1.83–1.74 (m, 1H), 1.65 (d, J = 1.9 Hz, 3H), 1.54 (dd, J = 2.8, 1.0 Hz, 3H), 1.53–1.46 (m, 1H), 0.92 (d, J = 6.9 Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ = 177.3, 138.7, 125.5, 59.9, 57.4, 51.7, 44.6, 37.8, 32.8, 32.0, 22.5, 20.7, 15.3; HRMS (ESI) m/z calcd for $\text{C}_{13}\text{H}_{23}\text{O}_3$ $[\text{M} + \text{H}]^+$: 227.1642, found: 227.1642.

Methyl (1R,2R)-2-methyl-1-(2-oxoethyl)-5-(propan-2-ylidene)cyclopentane-1-carboxylate (53): Alcohol **52** (3.35 g, 14.8 mmol, 1.0 eq.) was dissolved in dry CH_2Cl_2 (150 mL), DMP (7.05 g, 16.3 mmol, 1.1 eq.) was added and the reaction mixture was stirred for 2 h at rt. Sat. aq. NaHCO_3 solution (50 mL) was added and the aqueous layer was extracted with CH_2Cl_2 (5 x 50 mL). The combined organic layers were washed with brine (25 mL), dried over Na_2SO_4 , filtered and the solvent was removed under reduced pressure. The residue was purified by flash column chromatography (pentane/ Et_2O 4:1) to obtain the aldehyde **53** (2.95 g, 13.2 mmol, 89%) as colorless oil; R_f = 0.38 (pentane/ Et_2O 5:1); $[\alpha]_D^{25}$ = +23.6 (c 0.59, CHCl_3); FTIR (neat): $\tilde{\nu}$ = 2952, 2735, 2361, 1722, 1457, 1222, 1058, 892, 767 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ = 9.66 (t, J = 2.9 Hz, 1H), 3.68 (s, 3H), 2.93 (dd, J = 16.1, 2.8 Hz, 1H), 2.74 (dd, J = 16.1, 3.1 Hz, 1H), 2.56 (dd, J = 15.8, 7.5 Hz, 1H), 2.31–2.13 (m, 2H), 1.89–1.79 (m, 1H), 1.66 (d, J = 1.7 Hz, 3H), 1.61 (m, 1H), 1.56 (dd, J = 2.6, 1.1 Hz, 3H), 0.93 (d, J = 6.8 Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ = 202.5, 175.3, 137.4, 126.6, 56.2, 51.9, 48.3, 45.9, 32.6, 32.0, 22.6, 20.7, 14.6; HRMS (ESI) m/z calcd for $\text{C}_{13}\text{H}_{21}\text{O}_3$ $[\text{M} + \text{H}]^+$: 225.1485, found: 225.1485.

Methyl (1R,2R)-1-(4-ethoxy-2,4-dioxobutyl)-2-methyl-5-(propan-2-ylidene)cyclopentane-1-carboxylate (54): Ethyl diazoacetate (3.08 g, 26.4 mmol, 2.0 eq.) was dissolved in dry CH_2Cl_2 (25 mL), SnCl_2 (255 mg, 1.32 mmol, 0.1 eq.) was added and the reaction mixture was stirred at rt. A solution of the aldehyde **53** (2.95 g, 13.2 mmol, 1.0 eq.) in dry CH_2Cl_2 (30 mL) was added dropwise to the mixture. The color changed to yellow and formation of N_2 was observed. The reaction mixture was stirred for 12 h at rt, then it was diluted with CH_2Cl_2 (50 mL), filtered over celite and the solvent was removed under reduced pressure. The residue was purified by flash column chromatography (pentane/ Et_2O 7:1 to 2:1) to obtain the β -keto ester **54** (3.82 g, 12 mmol, 93%) as slightly orange oil; R_f = 0.24

(pentane/Et₂O 5:1); $[\alpha]_D^{25} = +49.6$ (*c* 0.75, CHCl₃); FTIR (neat): $\tilde{\nu} = 2950, 1717, 1644, 1454, 1306, 1224, 1183, 1155, 1033 \text{ cm}^{-1}$; (ratio keto ester/enol ester 10:1, only the data for the keto ester **54** is reported for the NMR set) ¹H NMR (400 MHz, CDCl₃) $\delta = 4.17$ (q, *J* = 7.1 Hz, 2H), 3.65 (d, *J* = 1.8 Hz, 3H), 3.41 (d, *J* = 5.5 Hz, 2H), 3.23 (d, *J* = 16.7 Hz, 1H), 2.95 (d, *J* = 16.7 Hz, 1H), 2.57–2.41 (m, 2H), 2.41–2.25 (m, 1H), 1.85–1.73 (m, 1H), 1.66–1.61 (m, 3H), 1.55–1.46 (m, 1H), 1.52 (d, *J* = 1.5 Hz, 3H), 1.26 (t, *J* = 7.1 Hz, 3H), 0.85 (d, *J* = 6.9 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) $\delta = 201.2, 175.8, 167.3, 138.5, 124.6, 61.4, 56.9, 51.7, 50.8, 46.6, 43.6, 32.8, 31.9, 22.5, 20.5, 14.7, 14.2$; HRMS (ESI) *m/z* calcd for C₁₇H₂₇O₅ [M + H]⁺: 311.1853, found: 311.1855.

6-Ethyl 3a-methyl (3R,3aR,6R)-6-chloro-3,7,7-trimethyl-5-oxo-2,3,4,5,6,7-hexahydro-3aH-indene-3a,6-dicarboxylate (55): β-Keto ester **54** (40.0 mg, 129 μmol, 1.0 eq.) was dissolved in degassed Ac₂O (3.0 mL, freeze-thaw method, 3 cycles). Flame dried LiCl (54.7 mg, 1.29 mmol, 10 eq.) was added and the reaction mixture was heated to 50 °C. Mn(OAc)₃ · 2 H₂O (141 mg, 516 μmol, 4.0 eq.) was added and the reaction mixture was stirred for 5 h at this temperature. Dist. water (5 mL) was added and the mixture was extracted with EtOAc (5 x 5 mL). The combined organic layers were washed with brine (5 mL), dried over Na₂SO₄, filtered and evaporated under reduced pressure. The residue was purified by flash column chromatography (pentane/Et₂O 6:1 to 3:1) to obtain the oxo-dicarboxylate **55** (17.1 mg, 50 μmol, 39%) as white solid; Mp: 66.2–67.8 °C; *R*_f = 0.26 (pentane/Et₂O 6:1); $[\alpha]_D^{25} = -53.2$ (*c* 0.73, CHCl₃); FTIR (neat): $\tilde{\nu} = 2982, 2931, 1728, 1459, 1408, 1385, 1366, 1346, 1246, 1225, 1105, 1034, 979, 894, 833, 757, 708, 606 \text{ cm}^{-1}$; ¹H NMR (400 MHz, CDCl₃) $\delta = 5.91\text{--}5.90$ (m, 1H), 4.36–4.21 (m, 2H), 3.68 (s, 3H), 3.49 (d, *J* = 15.3 Hz, 1H), 2.99 (d, *J* = 15.5 Hz, 1H), 2.55–2.42 (m, 2H), 2.35–2.23 (m, 1H), 1.38 (s, 3H), 1.32 (t, *J* = 7.1 Hz, 3H), 1.21 (s, 3H), 0.97 (d, *J* = 6.7 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) $\delta = 199.7, 173.9, 165.8, 146.1, 129.9, 79.0, 62.7, 60.2, 52.1, 50.8, 45.2, 43.6, 39.2, 24.1, 22.1, 14.2, 14.1$; HRMS (ESI) *m/z* calcd for C₁₇H₂₄O₅Cl [M + H]⁺: 343.1307, found: 343.1308.

6-Ethyl 3a-methyl (3R,3aR,5R,6R)-6-chloro-5-hydroxy-3,7,7-trimethyl-2,3,4,5,6,7-hexahydro-3aH-indene-3a,6-dicarboxylate (56): The oxo-dicarboxylate **55** (1.05 g,

3.05 mmol, 1.0 eq.) was dissolved in dry MeOH (50 mL). The solution was cooled to 0 °C and NaBH₄ (294 mg, 7.63 mmol, 2.5 eq.) was added in small portions. The reaction mixture was stirred at 0 °C for 1 h, then the cooling bath was removed and the reaction mixture was allowed to stir for 3 h at rt. The reaction mixture was cooled to 0 °C and quenched with brine (50 mL). The aqueous layer was extracted with EtOAc (5 x 50 mL) and the combined organic layers were washed with brine (30 mL). The organic layers were dried over Na₂SO₄, filtered and evaporated under reduced pressure. The residue was purified by flash column chromatography (pentane/Et₂O 6:1 to 3:1) to afford the hydroxy-dicarboxylate **56** (904 mg, 2.6 mmol, 86%) as white solid; Mp: 82.1–83.0 °C; R_f = 0.14 (pentane/Et₂O 6:1); [α]_D²⁶ = +42.4 (c 0.28, CHCl₃); FTIR (neat): $\tilde{\nu}$ = 3457, 2967, 2929, 2338, 1734, 1692, 1450, 1409, 1384, 1363, 1284, 1239, 1193, 1105, 1088, 1032, 995, 912, 887, 853, 819, 757, 702, 627 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ = 5.87 (dd, *J* = 3.2, 1.7 Hz, 1H), 4.50 (s, 1H), 4.36–4.21 (m, 3H), 3.69 (s, 3H), 2.88 (dd, *J* = 14.7, 2.8 Hz, 1H), 2.42 (ddd, *J* = 14.9, 7.4, 3.1 Hz, 1H), 2.32–2.20 (m, 1H), 2.17 (ddd, *J* = 14.9, 10.3, 1.7 Hz, 1H), 2.03 (dd, *J* = 14.8, 3.7 Hz, 1H), 1.53 (s, 3H), 1.38–1.31 (m, 6H), 0.97 (d, *J* = 6.9 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ = 176.6, 170.7, 147.6, 129.6, 75.0, 74.2, 62.2, 56.4, 51.7, 49.9, 42.3, 38.9, 35.8, 26.5, 26.1, 14.7, 14.1; HRMS (ESI) *m/z* calcd for C₁₇H₂₆O₅Cl [M + H]⁺: 345.1463, found: 345.1462; X-ray crystal structure is given in the supporting information.

9-Ethyl 5a-methyl (1R,4R,5aR,6R,9R)-9-chloro-1,6-dimethyl-1,2,4,5,6,7-hexahydro-5aH-1,4-methanocyclopenta[d]oxepine-5a,9-dicarboxylate (57) and 9-Ethyl 5a-methyl (1R,2R,4R,5aR,6R,9R)-2-acetoxy-9-chloro-1,6-dimethyl-1,2,4,5,6,7-hexahydro-5aH-1,4-methanocyclopenta[d]oxepine-5a,9-dicarboxylate (58): Hydroxy-dicarboxylate **56** (3.48 g, 10.1 mmol, 1.0 eq.) was dissolved in degassed cyclohexane (200 mL, freeze-thaw method, 3 cycles). Pb(IV) acetate (11.5 g, 25.2 mmol, 2.5 eq.), iodine (6.54 g, 25.2 mmol, 2.5 eq.) and CaCO₃ (4.04 g, 40.4 mmol, 4.0 eq.) were added. The reaction mixture was stirred at rt under irradiation of Hg-light (low pressure lamp) for three days and Pb(IV) acetate (45.7 g, 101 mmol, 10 eq.), iodine (26.2 g, 101 mmol, 10 eq.) and CaCO₃ (10.3 g, 101 mmol, 10 eq.) were added in four equal portions (each 2.5 eq.) during this time. The violet reaction mixture was cooled to 0 °C and sat. aq. Na₂S₂O₃ (400 mL)

was added until the mixture turned colorless. The mixture was extracted with EtOAc (4 x 200 mL), washed with brine (100 mL), dried over Na₂SO₄, filtered and evaporated under reduced pressure. The crude was purified by flash column chromatography (pentane/Et₂O 5:1 to 3:1) to afford the chloro-furan **57** (major product) (2.82 g, 8.2 mmol, 81%) as colorless solid and furan-acetate **58** (minor product) (600 mg, 1.5 mmol, 15%) as colorless solid; **57**: Mp: 96.5–97.1 °C; *R_f* = 0.16 (pentane/Et₂O 4:1); [α]_D²⁵ = –50.5 (*c* 0.37, CHCl₃); FTIR (neat): $\tilde{\nu}$ = 2961, 2945, 2906, 2360, 1736, 1721, 1453, 1383, 1228, 1081, 1031, 981, 936, 876, 772, 726 cm^{–1}; ¹H NMR (400 MHz, CDCl₃) δ = 5.83 (s, 1H), 4.71 (d, *J* = 4.8 Hz, 1H), 4.24 (q, *J* = 7.1 Hz, 1H), 3.85 (dd, *J* = 8.1, 0.8 Hz, 1H), 3.72–3.66 (m, 3H), 3.59 (d, *J* = 8.2 Hz, 1H), 3.07 (dd, *J* = 14.4, 4.9 Hz, 1H), 2.46–2.32 (m, 2H), 2.21–2.02 (m, 1H), 1.90 (d, *J* = 14.2 Hz, 1H), 1.35 (s, 3H), 1.29 (td, *J* = 7.1, 0.8 Hz, 3H), 0.91 (d, *J* = 6.6 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ = 173.8, 168.5, 145.6, 129.8, 79.6, 73.7, 73.6, 62.4, 57.5, 51.8, 49.5, 48.7, 39.4, 37.0, 15.1, 14.1, 13.4; HRMS (ESI) *m/z* calcd for C₁₇H₂₃O₅ClNa [M + Na]⁺: 365.1126, found: 365.1125; X-ray crystal structure is given in the supporting information; **58**: Mp: 130.5–131.5 °C; *R_f* = 0.31 (pentane/Et₂O 2:1); [α]_D²⁵ = +31.8 (*c* 0.57, CHCl₃); FTIR (neat): $\tilde{\nu}$ = 2959, 1755, 1724, 1440, 1377, 1255, 1218, 1178, 1078, 1019, 991, 934, 882, 776, 738 cm^{–1}; ¹H NMR (500 MHz, CDCl₃) δ = 6.04–6.02 (m, 1H), 5.97 (s, 1H), 5.15 (d, *J* = 4.5 Hz, 1H), 4.33–4.18 (m, 2H), 3.73 (s, 3H), 3.12 (dd, *J* = 14.4, 5.2 Hz, 1H), 2.46–2.37 (m, 2H), 2.18–2.09 (m, 1H), 1.97 (s, 3H), 1.90 (dd, *J* = 14.5, 0.8 Hz, 1H), 1.43 (s, 3H), 1.34 (t, *J* = 7.1 Hz, 3H), 0.94 (d, *J* = 6.6 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ = 173.4, 169.0, 168.0, 142.8, 132.9, 97.6, 80.1, 71.2, 62.4, 57.6, 54.0, 52.1, 48.5, 39.3, 36.0, 21.1, 15.2, 14.1, 12.5; HRMS (ESI) *m/z* calcd for C₁₉H₂₅O₇ClNa [M + Na]⁺: 423.1181, found: 423.1188; X-ray crystal structure is given in the supporting information.

Conversion of acetyl-furan **58** to chloro-furan **57**: Acetyl-furan **58** (702 mg, 1.75 mmol, 1.0 eq.) was dissolved in dry CH₂Cl₂ (120 mL) and cooled to –78 °C. Triethylsilane (514 mg, 4.38 mmol, 2.5 eq.) and boron trifluoride etherate (621 mg, 4.38 mmol, 2.5 eq.) were added. The reaction mixture was stirred 1 h at –78 °C and then was allowed to warm up to rt over a period of 12 h. The reaction mixture was cooled to 0 °C and dist. water (80 mL) was slowly added. The mixture was extracted with Et₂O (3 x 80 mL), washed with

brine (50 mL), dried over Na₂SO₄, filtered and evaporated under reduced pressure. The crude was purified by flash column chromatography (pentane/Et₂O 4:1) to afford the chloro-furan **57** (525 mg, 1.53 mmol, 88%) as colorless solid.

9-Ethyl 5a-methyl (1R,4R,5aR,6R,9S)-1,6-dimethyl-1,2,4,5,6,7-hexahydro-5aH-1,4-methanocyclopenta[d]oxepine-5a,9-dicarboxylate (59): Chloro-furan **57** (483 mg, 1.41 mmol, 1.0 eq.) was dissolved in dry benzene (35 mL). Sn(Bu)₃H (487 mg, 1.62 mmol, 1.15 eq.) and AIBN (23.9 mg, 0.14 mmol, 0.1 eq.) were added to the reaction mixture. The reaction mixture was heated to 100 °C and stirred for 2 h. The reaction was cooled to rt and then evaporated under reduced pressure. The residue was purified by flash column chromatography (pentane/Et₂O 3:1) to afford the hydro-furan **59** (411 mg, 1.34 mmol, 95%) as colorless oil; *R*_f = 0.16 (pentane/Et₂O 3:1); [α]_D²⁵ = +9.0 (*c* 0.78, CHCl₃); FTIR (neat) $\tilde{\nu}$ = 2955, 2880, 1725, 1450, 1378, 1337, 1196, 1155, 1049, 1023, 989, 925, 664, 627 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ = 5.72 (s, 1H), 4.56 (t, *J* = 5.2 Hz, 1H), 4.20–4.04 (m, 2H), 3.78 (d, *J* = 7.7 Hz, 1H), 3.69 (s, 3H), 3.46 (d, *J* = 7.7 Hz, 1H), 3.01 (dd, *J* = 14.2, 4.7 Hz, 1H), 2.83 (dd, *J* = 5.9, 1.1 Hz, 1H), 2.39–2.23 (m, 2H), 2.13–2.03 (m, 1H), 1.71 (d, *J* = 14.5 Hz, 1H), 1.35 (s, 3H), 1.23 (t, *J* = 7.1 Hz, 3H), 0.90 (d, *J* = 6.6 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ = 174.3, 169.7, 145.8, 126.6, 77.4, 76.7, 60.3, 58.3, 55.7, 51.7, 48.9, 44.8, 39.0, 37.7, 17.4, 15.2, 14.4; HRMS (ESI) *m/z* calcd for C₁₇H₂₅O₅ [M + H]⁺: 309.1697, found: 309.1698.

6-Ethyl 3a-methyl (3R,3aR,6S,7R)-5-acetoxy-7-(acetoxymethyl)-3,7-dimethyl-2,3,4,5,6,7-hexahydro-3aH-indene-3a,6-dicarboxylate (60) and 6-Ethyl 3a-methyl (3R,3aS,7R)-7-(acetoxymethyl)-3,7-dimethyl-2,3,6,7-tetrahydro-3aH-indene-3a,6-dicarboxylate (61): Hydro-furan **59** (44.7 mg, 145 μ mol, 1.0 eq.) was dissolved in Ac₂O (0.9 mL). The solution was cooled to –20 °C and boron trifluoride etherate (56.2 μ L, 435 μ mol, 3.0 eq.) was added. The reaction was allowed to warm up to rt over 12 h. The reaction mixture was cooled to 0 °C and dist. water (1.0 mL) was added dropwise. The reaction mixture was stirred for 10 min and warmed up to rt. The mixture was extracted with Et₂O (3 x 1.5 mL), washed with brine (1.5 mL), dried over Na₂SO₄, filtered and evaporated under reduced pressure. The residue was purified by flash column

chromatography (pentane/Et₂O 2:1) to afford the diacetylated compound **60** (35.4 mg, 87 μ mol, 60%) as a yellow oil and the monoacetylated compound **61** (31.7 mg, 46 μ mol, 32%) as colorless oil; **60**: R_f = 0.17 (pentane/Et₂O 2:1); $[\alpha]_D^{25}$ = +8.9 (*c* 0.34, CHCl₃); FTIR (neat): $\tilde{\nu}$ = 3480, 2956, 1727, 1440, 1367, 1224, 1028, 908, 790, 664 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ = 5.83 (s, 1H), 5.30–5.23 (m, 1H), 4.19–4.03 (m, 2H), 3.77 (d, *J* = 11.3 Hz, 1H), 3.70 (d, *J* = 11.4 Hz, 1H), 3.65 (s, 3H), 3.13 (d, *J* = 5.4 Hz, 1H), 2.65 (dd, *J* = 12.3, 4.8 Hz, 1H), 2.51–2.36 (m, 2H), 2.26–2.14 (m, 1H), 2.07 (s, 3H), 2.03–1.97 (m, 1H), 1.99 (s, 3H), 1.23 (t, *J* = 7.1 Hz, 3H), 1.15 (s, 3H), 0.94 (d, *J* = 6.4 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ = 174.3, 170.9, 170.8, 169.9, 143.3, 131.1, 68.8, 68.7, 60.3, 59.5, 51.8, 49.5, 49.1, 40.5, 39.1, 34.5, 21.5, 21.1, 21.0, 14.5; HRMS (ESI) *m/z* calcd for C₂₁H₃₀O₈Na [M + Na]⁺: 433.1833, found: 433.1835; **61**: R_f = 0.32 (pentane/Et₂O 2:1); $[\alpha]_D^{26}$ = +223.8 (*c* 0.14, CHCl₃); FTIR (neat): $\tilde{\nu}$ = 2957, 1725, 1454, 1372, 1320, 1222, 1141, 1033, 793 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ = 6.22 (d, *J* = 10.0 Hz, 1H), 5.82 (s, 1H), 5.58 (dd, *J* = 10.0, 5.1 Hz, 1H), 4.14–4.01 (m, 2H), 3.92 (d, *J* = 10.9 Hz, 1H), 3.74 (d, *J* = 10.9 Hz, 1H), 3.65 (s, 3H), 3.15 (d, *J* = 5.1 Hz, 1H), 2.48–2.29 (m, 3H), 2.04 (s, 3H), 1.22 (t, *J* = 7.1 Hz, 3H), 1.19 (s, 3H), 1.03 (d, *J* = 6.3 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ = 173.1, 171.3, 170.9, 143.3, 132.3, 129.8, 122.1, 68.3, 60.8, 59.3, 51.8, 50.2, 47.9, 39.7, 38.8, 21.0, 19.4, 14.4, 14.3. HRMS (ESI) *m/z* calcd for C₁₉H₂₇O₆ [M + H]⁺: 351.1802, found: 351.1800.

6-Ethyl 3a-methyl (3R,3aR,6S,7R)-5-hydroxy-7-(hydroxymethyl)-3,7-dimethyl-2,3,4,5,6,7-hexahydro-3aH-indene-3a,6-dicarboxylate (62): The diacetylated product **60** (38.7 mg, 94.3 μ mol, 1.0 eq.) was dissolved in dry MeOH (1.5 mL), K₂CO₃ (53.2 mg, 0.38 mmol, 4.0 eq.) was added and the reaction mixture was stirred at rt for 16 h. The reaction mixture was diluted with EtOAc (5.0 mL), aq. HCl (1 M, 2.0 mL) was added and the mixture was extracted with EtOAc (3 x 2.0 mL). The combined organic layers were washed with brine (1.5 mL), dried over Na₂SO₄, filtered and the solvent was removed under reduced pressure. The residue was purified by flash column chromatography (pentane/EtOAc 1:1) to afford the diol **62** (30.1 mg, 92 μ mol, 98%) as colorless sticky oil; R_f = 0.45 (pentane/EtOAc 1:1); $[\alpha]_D^{25}$ = -1.9 (*c* 0.25, CHCl₃); FTIR (neat): $\tilde{\nu}$ = 3448, 2954, 1721, 1448, 1178, 1027, 885, 780, 710, 660 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ =

5.86 (s, 1H), 4.20–4.04 (m, 3H), 3.70 (s, 3H), 3.20 (d, $J = 11.3$ Hz, 1H), 3.07 (d, $J = 11.3$ Hz, 1H), 2.70–2.60 (m, 2H), 2.51–2.40 (m, 2H), 2.22–1.88 (m, 4H), 1.24 (t, $J = 7.1$ Hz, 3H), 1.17 (s, 3H), 0.96 (d, $J = 6.7$ Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) $\delta = 175.4, 171.5, 142.5, 132.8, 66.6, 66.3, 60.4, 59.9, 54.3, 52.2, 48.6, 43.0, 39.0, 37.8, 21.4, 14.7, 14.4$; HRMS (ESI) m/z calcd for $\text{C}_{17}\text{H}_{26}\text{O}_6\text{Na}$ $[\text{M} + \text{Na}]^+$: 349.1622, found: 349.1626.

6-Ethyl 3a-methyl (3R,3aR,7S)-7-formyl-5-hydroxy-3,7-dimethyl-2,3,4,7-tetrahydro-3aH-indene-3a,6-dicarboxylate (63): To a flask charged with flame dried molecular sieves (4 Å, 120 mg) was added a solution of the diol **62** (58.1 mg, 178 μmol , 1.0 eq.) in dry CH_2Cl_2 (5.0 mL). NMO (56.0 mg, 0.46 mmol, 2.6 eq.) was added at rt and the reaction mixture was stirred for 20 min. TPAP (12.9 mg, 35 μmol , 0.2 eq.) was added and the solution was stirred at rt for 24 h, while the color turned from green to dark blue. The reaction mixture was filtered over celite and the solvent was removed under reduced pressure. The residue was purified by flash column chromatography (pentane/ Et_2O 3:1) to yield the enol **63** (39.0 mg, 0.12 mmol, 68%) as colorless oil; $R_f = 0.39$ (pentane/ Et_2O 3:1); $[\alpha]_D^{25} = -124.9$ (c 0.61, CHCl_3); FTIR (neat): $\tilde{\nu} = 2960, 2934, 2848, 1728, 1642, 1607, 1458, 1402, 1297, 1228, 1076, 863, 732, 631\text{ cm}^{-1}$; ^1H NMR (400 MHz, CDCl_3) $\delta = 12.94$ (s, 1H), 9.13 (s, 1H), 5.98 (s, 1H), 4.27–4.12 (m, 2H), 3.63 (s, 3H), 3.25 (d, $J = 16.4$ Hz, 1H), 2.54 (ddd, $J = 15.3, 7.3, 3.1$ Hz, 1H), 2.37–2.19 (m, 2H), 2.16 (d, $J = 16.3$ Hz, 1H), 1.47 (s, 3H), 1.23 (t, $J = 7.1$ Hz, 3H), 1.02 (d, $J = 6.9$ Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) $\delta = 197.3, 175.8, 172.8, 171.5, 141.3, 133.3, 99.0, 61.1, 60.2, 51.9, 50.8, 46.4, 39.8, 38.9, 22.7, 15.0, 14.1$; HRMS (ESI) m/z calcd for $\text{C}_{17}\text{H}_{23}\text{O}_6$ $[\text{M} + \text{H}]^+$: 323.1489, found: 323.1490.

6-Ethyl 3a-methyl (3R,3aR,7S)-7-formyl-6-hydroxy-3,7-dimethyl-5-oxo-2,3,4,5,6,7-hexahydro-3aH-indene-3a,6-dicarboxylate (64): The enol **63** (89.0 mg, 276 μmol , 1.0 eq.) was dissolved in dry *i*-PrOH (5 mL), then degassed (freeze-thaw method, 2 cycles) and refilled with oxygen. $\text{CeCl}_3 \cdot 7\text{H}_2\text{O}$ (105 mg, 276 μmol , 1.0 eq.) was added at rt and oxygen was bubbled through the colorless solution for 15 min. The solution was stirred under oxygen atmosphere at rt for 20 h. Dist. water (5 mL) was added, the mixture was extracted with EtOAc (3 x 5 mL), the combined organic layers were washed with brine (3

mL), dried over Na₂SO₄, filtered and the solvent was removed under reduced pressure. The crude was purified by flash column chromatography (pentane/Et₂O 2:1) to afford the α -hydroxylated product **64** (40 mg, 119 μ mol, 43%) as colorless oil; R_f = 0.37 (pentane/Et₂O 1:1); $[\alpha]_D^{25}$ = -68.8 (c 0.53, CHCl₃); FTIR (neat): $\tilde{\nu}$ = 3454, 2959, 1724, 1453, 1373, 1217, 1162, 1099, 1001, 916, 859, 732, 653 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ = 9.40 (s, 1H), 6.11 (s, 1H), 4.31 (q, J = 7.1 Hz, 2H), 3.91 (s, 1H), 3.64 (s, 3H), 3.51 (d, J = 14.7 Hz, 1H), 2.84 (d, J = 14.7 Hz, 1H), 2.59–2.45 (m, 2H), 2.32–2.20 (m, 1H), 1.40 (s, 3H), 1.28 (t, J = 7.2 Hz, 3H), 0.98 (d, J = 6.7 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ = 202.8, 197.3, 172.5, 170.4, 138.8, 135.1, 82.9, 63.8, 61.1, 58.7, 52.2, 50.6, 44.4, 39.3, 14.6, 14.5, 14.1; HRMS (ESI) m/z calcd for C₁₇H₂₃O₇ [M + H]⁺: 339.1438, found: 339.1441.

(3*aR*,4*R*,6*aR*,7*R*,9*bR*)-3*a*-Hydroxy-7,9*b*-dimethyl-1,3*a*,4,7,8,9*b*-hexahydro-3*H*,6*H*-4,6*a*-methanocyclopenta[*c*]furo[3,4-*e*]oxepine-3,6-dione (**65**): The α -hydroxylated ketone **64** (21.3 mg, 63.0 μ mol, 1.0 eq.) was dissolved in dry MeOH (1.0 mL) and cooled to 0 °C. NaBH₄ (11.9 mg, 315 μ mol, 5.0 eq.) was added in small portions and the reaction mixture was allowed to warm up to rt over 11 h. The reaction mixture was cooled to 0 °C and dist. water (1.5 mL) was added. The aq. layer was extracted with EtOAc (4 x 1.5 mL), dried over Na₂SO₄, filtered and the solvent was removed under reduced pressure to afford the crude diol as colorless solid. The crude diol (21 mg, 70.9 μ mol, 1.0 eq.) was dissolved in dry toluene (1.0 mL) and *p*-toluensulfonic acid monohydrate (2.7 mg, 14.2 μ mol, 0.2 eq.) was added. The reaction mixture was stirred at 77 °C for 2 h. The reaction mixture was diluted with EtOAc (3 mL) and washed with aq. sat. NaHCO₃ solution (2.0 mL). The aq. layer was extracted with EtOAc (3 x 1.5 mL), the combined organic layers was washed with brine (1.5 mL), dried over Na₂SO₄, filtered and the solvent was removed under reduced pressure. The residue was purified by flash column chromatography (pentane/EtOAc 3:1) to afford the dilactone **65** (4.4 mg, 17 μ mol, 27%, over two steps) as white solid; Mp: 129.5–130.2 °C; R_f = 0.53 (pentane/EtOAc 2:1); $[\alpha]_D^{25}$ = +75.0 (c 0.47, CHCl₃); FTIR (neat): $\tilde{\nu}$ = 3448, 2921, 2851, 1782, 1753, 1457, 1376, 1326, 1221, 1185, 1113, 1016, 1000, 938, 818, 732, 639 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ = 5.87 (s, 1H), 4.63 (d, J = 5.9 Hz, 1H), 3.98 (q, J = 9.5 Hz, 2H), 2.58 (ddd, J = 15.6, 8.1, 2.9 Hz, 1H),

2.48 (dd, $J = 11.9, 5.9$ Hz, 1H), 2.44–2.37 (m, 1H), 2.32 (ddd, $J = 15.5, 9.7, 1.9$ Hz, 1H), 2.21 (d, $J = 12.1$ Hz, 1H), 1.36 (s, 3H), 1.32 (d, $J = 6.9$ Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) $\delta = 177.7, 174.9, 140.6, 129.8, 78.9, 78.0, 74.4, 56.6, 43.3, 39.2, 37.0, 35.0, 21.4, 13.3$; HRMS (ESI) m/z calcd for $\text{C}_{14}\text{H}_{16}\text{O}_5\text{Na}$ $[\text{M} + \text{Na}]^+$: 287.0890, found: 287.0894.

Supporting Information

Reaction conditions for the oxidation of compound **33**. ^1H - and ^{13}C -NMR spectra. X-Ray crystal structure analysis data.

Acknowledgments

We thank Dr. Heinz Nadig for HRMS measurements and acknowledge the Swiss National Science Foundation for financial support (200020_163151 and 200021_144028).

References

-
- (1) (a) Xu, J.; Lacoske, M. H.; Theodorakis, E. A. *Angew. Chem. Int. Ed.* **2014**, *53*, 956. (b) Akagi, M.; Matsui, N.; Akae, H.; Hirashima, N.; Fukuishi, N.; Fukuyama, Y.; Akagi, R. *Journal of Pharmacological Sciences* **2015**, *127*, 155.
- (2) (a) Birman, V. B.; Danishefsky, S. J. *J. Am. Chem. Soc.* **2002**, *124*, 2080. (b) Ding, R.; Sun, B.-F.; Lin, G.-Q. *Org. Lett.* **2012**, *14*, 4446. (c) Koshiba, T.; Yokoshima, S.; Fukuyama, T. *Org. Lett.* **2009**, *11*, 5354. (d) Watanabe, H.; Takano, M.; Umino, A.; Ito, T.; Ishikawa, H.; Nakada, M. *Org. Lett.* **2007**, *9*, 359. (e) Ogura, A.; Yamada, K.; Yokoshima, S.; Fukuyama, T. *Org. Lett.* **2012**, *14*, 1632. (f) Elamparuthi, E.; Fellay, C.; Neuburger, M.; Gademann, K. *Angew. Chem. Int. Ed.* **2012**, *51*, 4071. (g) Burch, P.;

-
- Binaghi, M.; Scherer, M.; Wetzel, C.; Bossert, D.; Eberhardt, L.; Neuburger, M.; Scheiffele, P.; Gademann, K. *Chem. Eur. J.* **2013**, *19*, 2589.
- (3) (a) Yang, C.-S.; Kuono, I.; Kawano, N.; Sato, S. *Tetrahedron Lett.* **1988**, *29*, 1165. (b) Fukuyama, Y.; Huang, J.-M., *Studies in Natural Products Chemistry*; A. Rahman, Ed.; Elsevier: New York, **2005**, Vol. 32, p 395.
- (4) (a) Carcache, D. A.; Cho, Y. S.; Hua, Z.; Tian, Y.; Li, Y.-M.; Danishefsky, S. J. *J. Am. Chem. Soc.* **2006**, *128*, 1016. (b) Trzoss, L.; Xu, J.; Lacoske, M. H.; Mobley, W. C.; Theodorakis, E. A. *Chem. Eur. J.* **2013**, *19*, 6398.
- (5) Yokoyama, R.; Huang, J.-M.; Yang, C.-S.; Fukuyama, Y. *J. Nat. Prod.* **2002**, *65*, 527.
- (6) Kubo, M.; Okada, C.; Huang, J.-M.; Harada, K.; Hioki, H.; Fukuyama, Y. *Org. Lett.* **2009**, *11*, 5190.
- (7) Kubo, M.; Kobayashi, K.; Huang, J.-M.; Harada, K.; Fukuyama, Y. *Tetrahedron Lett.* **2012**, *53*, 1231.
- (8) (a) Cho, Y. S.; Carache, D. A.; Tian, Y.; Li, Y.-M.; Danishefsky, S. J. *J. Am. Chem. Soc.* **2004**, *126*, 14358. (b) Yang, Y.; Fu, X.; Chen, J.; Zhai, H. *Angew. Chem. Int. Ed.* **2012**, *51*, 9825. (c) Trzoss, L.; Xu, J.; Lacoske, M. H.; Mobley, W.C.; Theodorakis, E. A. *Org. Lett.* **2011**, *13*, 4554. (d) Chen, X.; Micalizio, G. C. *J. Am. Chem. Soc.* **2016**, *138*, 1150. (e) Harada, K.; Imai, A.; Uto, K.; Carter, R. G.; Kubo, M.; Hioli, H.; Fukuyama, Y. *Tetrahedron* **2015**, *71*, 2199.
- (9) (a) Xu, J.; Trzoss, L.; Chang, W. K.; Theodorakis, E. A. *Angew. Chem. Int. Ed.* **2011**, *50*, 3672. (b) Siler, D. A.; Mighion, J. D.; Sorensen, E. J. *Angew. Chem. Int. Ed.* **2014**, *53*, 5332. (c) Paterson, I.; Xuan, M.; Dalby, S. M. *Angew. Chem. Int. Ed.* **2014**, *53*, 7286. (d) Shen, Y.; Li, L.; Pan, Z.; Wang, Y.; Li, J.; Wang, K.; Wang, X.; Zhang, Y. Hu, T.; Zhang, Y. *Org. Lett.* **2015**, *17*, 5480.
- (10) Lu, H.-H.; Martinez, M. D.; Shenvi, R. A. *Nature Chemistry* **2015**, *7*, 604.
- (11) Gomes, J. *Synthesis of Majucin-Type Sesquiterpenes and Immobilization and Visualization of Quorum Sensing Signaling Molecules*, PhD-Thesis, University of Basel, **2014**. Doi: 10.5451/unibas-006288268, published Sept. 3, 2014.
- (12) (a) Wolinsky, J.; Chan, D. *J. Org. Chem.* **1965**, *30*, 41. (b) Coates, R. M.; Vettel, P. *R. J. Org. Chem.* **1980**, *45*, 5430.

-
- (13) (a) Miller, J. J.; Sigman, M. S. *J. Am. Chem. Soc.* **2007**, *129*, 2752. (b) Knochel, P.; Rao, C. J. *Tetrahedron* **1993**, *49*, 29. (c) Angell, R.; Parsons, P. J.; Naylor, A. *Synlett* **1993**, 189. (d) Suzuki, K.; Takayama, H. *Org. Lett.* **2006**, *8*, 4605. (e) Takai, E.; Kimura, K.; Kuroda, T.; Hiyama, T.; Nozaki, H. *Tetrahedron Lett.* **1983**, *24*, 5281.
- (14) Janini, T. E. Sampson, P. *J. Org. Chem.* **1997**, *62*, 5069.
- (15) Niwa, H.; Nisiwaki, M.; Tsukada, I.; Ishigaki, T.; Ito, S.; Wakamatsu, K.; Mori, T.; Ikagawa, M.; Yamada, K. *J. Am. Chem. Soc.* **1990**, *112*, 9001.
- (16) Ramachary, D. P.; Kishor, M. *Org. Biomol. Chem.* **2008**, *6*, 4176.
- (17) Nicolaou, K. C.; Tang, Y.; Wang, J.; Stepan, A. F.; Li, A.; Montero, A. *J. Am. Chem. Soc.* **2007**, *129*, 14850.
- (18) (a) Alexakis, A.; Berlan, J.; Besace, Y. *Tetrahedron Lett.* **1986**, *27*, 1047. (b) Lipshutz, B. H. *Tetrahedron Lett.* **1983**, *24*, 127.
- (19) Roberts, R. A.; Schüll, V.; Paquette, L. A., *J. Org. Chem.* **1983**, *48*, 2076.
- (20) (a) Furukawa, J.; Kawabata, R.; Nishimura, J. *Tetrahedron Lett.* **1966**, *28*, 3353. (b) Simmons, H. E.; Smith, R. D. *J. Am. Chem. Soc.* **1959**, *81*, 4256.
- (21) Lebel, H.; Marcoux, J.-F.; Molinaro, C.; Charette, A. B., *Chem. Rev.* **2003**, *103*, 977.
- (22) Gurjar, M. K.; Cherian, J.; Ramana, C. V. *Org. Lett.* **2004**, *6*, 317.
- (23) Pettigrew, J. D.; Paquette, L. A. *Heterocycles* **2010**, *80*, 99.
- (24) Lightner, D. A.; Gurst, J. E. *Organic Conformational Analysis and Stereochemistry from Circular Dichroism Spectroscopy*; Wiley-VCH: New York, **2000**, 173.
- (25) Kuroiwa, Y.; Matsumura, S.; Toshima, K, *Synlett* **2008**, *16*, 2523.
- (26) Dowd, P.; Zhang, W. *Tetrahedron Lett.* **1993**, *34*, 2095.
- (27) (a) Girard, P.; Namy, J. L.; Kagan, H. B. *J. Am. Chem. Soc.* **1980**, *102*, 2693. (b) Molander, G. A.; McKie, J. A. *J. Org. Chem.* **1993**, *58*, 7216.
- (28) Brockson, T. J.; Coelho, F.; Deprés, J.-P.; Greene, A. E.; Freire de Lima, M. E.; Hamelin, O.; Hartmann, B.; Kanazawa, A. M.; Wang, Y. *J. Am. Chem. Soc.* **2002**, *124*, 15313.
- (29) see supporting information for detailed conditions.
- (30) Premraj, R.; McLeod, M. D.; Simpson, G. W.; Banwell, M. G. *Heterocycles* **2012**, *85*, 2949.

-
- (31) FIRMENICH SA; BIRKBECK, Anthony A. Patent: WO2013/1027 A1, **2013**.
- (32) (a) Itoh, A.; Ozawa, S.; Oshima, K.; Nozaki, H. *Tetrahedron Lett.* **1980**, *21*, 361. (b) Itoh, A.; Ozawa, S.; Oshima, K.; Nozaki, H. *Bull. Chem. Soc. Jpn.* **1981**, *54*, 274.
- (33) (a) Danishefsky, J. D.; Armistead, D. M.; Wincott, F. E.; Selnick, H. G.; Hungate, R. *J. Am. Chem. Soc.* **1989**, *111*, 2967. (b) Danishefsky, J. D.; Armistead, D. M. *Tetrahedron Lett.* **1987**, *28*, 4959.
- (34) (a) Hosomi, A.; Sakurai, H. *Tetrahedron Letters* **1976**, *17*, 1295. (b) Hosomi, A.; Sakurai, H. *Tetrahedron Lett.* **1976**, *17*, 1295.
- (35) The stereogenic center at C-4 could not be assigned using NMR-techniques, see supporting informations for structural proposal
- (36) Daeppen, C.; Kaiser, M.; Neuburger, M.; Gademann, K. *Org. Lett.* **2015**, *17*, 5420.
- (37) (a) Holmquist, C. R.; Roskmap, E. J. *J. Org. Chem.* **1989**, *54*, 3258. (b) Holmquist, C. R.; Roskmap, E. *Tetrahedron Lett.* **1992**, *33*, 1131.
- (38) Cainelli, G.; Mihailovic, M. Lj.; Arigoni, D.; Jeger, O. *Helv. Chim. Acta* **1959**, *42*, 1124.
- (39) Kalvoda, J.; Heusler, K. *Synthesis* **1971**, 501.
- (40) Beszant, S.; Giannini, E.; Zanoni, G.; Vidari, G. *Tetrahedron: Asymmetry* **2002**, *13*, 1245.
- (41) Desroches, C.; Lopes, C.; Kessler, V.; Parola, S. *Dalton Trans.* **2003**, 2085.
- (42) (a) Swallen, L. C.; Boord, C. E. *J. Am. Chem. Soc.* **1930**, *52*, 651. (b) Numazawa, M.; Yamada, K. *Steroids* **1999**, *64*, 320.
- (43) Shoppee, C. W.; Coll, J. C.; Lack, R. E. *J. Chem. Soc. C* **1970**, 1893.
- (44) Christoffers, J.; Werner, T. *Synlett* **2002**, *1*, 119.
- (45) Bruker Analytical X-ray Systems, Inc. **2006**. Apex2, Version 2 User Manual, M86-E01078, Madison, WI.
- (46) Altomare, A.; Cascarano, G.; Giacovazzo, C.; Guagliardi, A.; Burla, M. C.; Polidori, G.; Camalli, M. *J. Appl. Cryst.* **1994**, *27*, 435.
- (47) Betteridge, P. W.; Carruthers, J. R.; Cooper, R. I.; Prout, K.; Watkin, D. J. *J. Appl. Cryst.* **2003**, *36*, 1487.
- (48) Szostak, M.; Spain, M.; Procter, D. J. *J. Org. Chem.* **2012**, *77*, 3049.